arXiv:1302.5760v1 [math.PR] 23 Feb 2013

WEAKLY ASYMMETRIC NON-SIMPLE EXCLUSION PROCESS AND THE KPZ EQUATION

*AMIR DEMBO AND †LI-CHENG TSAI

Abstract. We analyze a class of non-simple exclusion processes and the corresponding growth models by generalizing the Gärtner transformation. While the original argument, which matches three identities with three parameters, applies only to simple exclusion process, our approach is to identify the major non-linear drift term of the microscopic dynamical equation and to convert the drift term into a quasi-linear second order differential, yielding the stochastic heat equation (SHE) in the continuum limit. Using the generalized transformation, we prove convergence toward the KPZ equation, which is the first *universality* result of this kind in the context of KPZ universality class. While this class of exclusion processes are not explicitly solvable, we obtain the exact one-point limit distribution for the step and step Bernoulli initial conditions by using the previous results of [1, 7] and our convergence result.

1. Introduction

There is much interest in study of randomly growing interfaces, whose macroscopic behavior are usually described by non-linear partial differential equations. To describe the random growth mechanism, a noise term is introduced to the equation. A paradigm of such modeling equations is the KPZ equation [15], which in one space dimension reads

$$\partial_T \mathcal{H} = \frac{1}{2}\partial_X^2 \mathcal{H} - \frac{1}{2}(\partial_X \mathcal{H})^2 + \dot{W}, \quad \mathcal{H} = \mathcal{H}_T(X), \tag{1.1}$$

where $\dot{W}$ is the space-time white noise: $\mathbb{E}(\dot{W}_T(X)\dot{W}_S(Y)) = \delta(T-S)\delta(X-Y)$. Employing the non-rigorous renormalization argument of [8], Kardar, Parisi, and Zhang [15] showed that the scaling exponents of the fluctuation of $\mathcal{H}$, space X, and time T follow a $1:2:3$ ratio, which signifies a new universality class —the KPZ universality class. This universality class describes various phenomena including paper wetting, crack formation, and burning fronts (see [6, Section 1.1.2] and the reference therein). Moreover, it connects various models describing other phenomenon including last passage percolation, directed polymer in a random media, and polynuclear growth. Since [15], there has been intensive mathematical research on instances of explicitly solvable models in this universality class [4, 13, 14, 17]. See also [5, 6] and the reference therein. They all confirm the $1:2:3$ scaling exponents and have limit one-point statistics which are the GUE or GOE Tracy-Widom distribution. Exact one-point statistics of the KPZ equation are proven by [1, 7], and independently obtained by [18] through non-rigorous arguments. These distributions converge to the GUE Tracy-Widom distribution (or a variant of) as $T \to \infty$ under the $1:2:3$ scaling.

The quadratic term in (1.1) plagues rigorous mathematical treatment. Indeed, generic solutions to stochastic differential equations driven by white noise exhibit non-differentiability. Hence, $(\partial_X \mathcal{H})^2$ alone does not have a mathematical meaning. Rather, the correct mathematical interpretation is the Hopf-Cole transformation to the stochastic heat equation (SHE):

$$\mathcal{H}_T(X) = -\log \mathcal{Z}_T(X), \tag{1.2}$$

Date: February 26, 2013.

2010 *Mathematics Subject Classification.* Primary 60K35; Secondary 60H15, 82C22.

Key words and phrases. Kardar-Parisi-Zhang equation, stochastic heat equation, universality, asymmetric exclusion process.

* Research partially supported by NSF grant DMS-1106627.



Figure 1. Non-simple exclusion process.

$$\partial_T \mathcal{Z} = \frac{1}{2}\partial_X^2 \mathcal{Z} + \dot{W}\mathcal{Z}. \tag{1.3}$$

Since the SHE is linear, traditional stochastic calculus applies. Further, formally $\mathcal{H}_T(X)$ of (1.2) satisfies (1.1), so we *define* (1.1) by (1.2) and (1.3). This Hope-Cole transformation dates back to [12, 15]. A more comprehensive approach of defining (1.1) can be found in [11].

As we shall see, the Hopf-Cole solution of (1.1) also captures the asymptotic fluctuations of certain exclusion processes. Specifically, let $p_1, \ldots, p_m$, $q_1, \ldots, q_m$ be non-negative numbers adding to one: $\sum_k (p_k + q_k) = 1$. Consider finite range exclusion processes, where particles on the half integer lattice $\frac{1}{2} + \mathbb{Z}$ wait an exponential time with parameter one and then attempt to jump k units to the right with probability p_k, k units to the left with probability q_k, under the constraint that a jump is prohibited when the destination site is already occupied. The case $m = 1$ is called simple exclusion process. Let $\eta_t(x)$ be the occupation variable in spin convention at position $x - \frac{1}{2}$ and time $t \in [0, \infty)$

$$\eta_t(x) := \begin{cases} \ 1 \text{ , when the site } x - \frac{1}{2} \text{ is occupied at time } t, \\ -1, \text{ otherwise.} \end{cases}$$

We associate a height function h by doing discrete integration of η, that is

$$h_t(x) = -N(t) + \begin{cases} \sum_{0<y\le x} \eta_t(y) \text{ , when } x \ge 0, \\ -\sum_{x<y\le 0} \eta_t(y), \text{ when } x < 0, \end{cases} \tag{1.4}$$

where $N(t)$ is the net flow of particles through $x = 0$ during the time interval $[0, t]$, counting right moving particles as positive. Note that the discrete gradient of h yields η:

$$h_t(x) - h_t(x-1) = \eta_t(x).$$

The height h represents the interface of a growth with the following dynamics: when a particle jumps k unit to the left(right), it adds(removes) a box at each of the k sites along the way cf. Figure 1.

For weakly asymmetric *simple* exclusion process, Gärtner [10] showed that the following discrete version of the Hopf-Cole transformation:

$$H_T^\varepsilon(X) := \lambda_\varepsilon h\left(\frac{X}{\varepsilon}, \frac{T}{\gamma_\varepsilon \varepsilon^{3/2}}\right) - \frac{\nu_\varepsilon T}{\gamma_\varepsilon \varepsilon^{3/2}}, \quad Z_T^\varepsilon(X) := \exp\left(-H_T^\varepsilon(X)\right), \tag{1.5}$$

where

$$\gamma_\varepsilon := q - p := \varepsilon^{\frac{1}{2}}, \tag{1.6a}$$

$$\lambda_\varepsilon := \frac{1}{2}\log\left(\frac{q}{p}\right) = \frac{1}{2}\log\left(\frac{1+\varepsilon^{\frac{1}{2}}}{1-\varepsilon^{\frac{1}{2}}}\right) = \varepsilon^{\frac{1}{2}} + \frac{1}{3}\varepsilon^{\frac{3}{2}} + O(\varepsilon^{\frac{5}{2}}), \tag{1.6b}$$

$$\nu_\varepsilon := \alpha\Big(1-(1-\varepsilon)^{\frac{1}{2}}\Big) = \alpha\Big(\frac{1}{2}\varepsilon + \frac{1}{8}\varepsilon^2 + O(\varepsilon^3)\Big), \quad \alpha = 1, \tag{1.6c}$$

converts the dynamics of h into a discretization of the SHE. This transformation linearizes the drift term of the microscopic dynamics equation by matching the four identities corresponding to $(\eta_t(x), \eta_t(x+1)) \in \{\pm 1\}^2$ (which actually degenerates into three identities) using a suitable choice of the parameters γ, λ, and ν. Going beyond simple exclusion process ($m > 1$), we encounter $2^m - 1 > 3$ identities, so this type of arguments fail. Nevertheless, for the class of exclusion processes with jump rates satisfying

$$\begin{aligned} &p_k = r_k p, \ q_k = r_k q, \\ &\text{where } p, q, r_1, \dots, r_m \in [0,1], \quad q - p = \gamma_\varepsilon, \ \sum_{k=1}^{m} r_k = 1, \end{aligned} \tag{1.7}$$

we show that the choice (1.6) of parameters with $\alpha = 1$ replaced by

$$\alpha := \sum_{k=1}^{m} k^2 r_k \tag{1.8}$$

eliminates the *major* non-linearity, leaving a drift term that behaves like a discrete Laplacian.

Without lost of generality we may assume that the associated (free) random walk on $\frac{1}{2} + \mathbb{Z}$ is irreducible, that is

$$\gcd\{k : r_k \neq 0\} = 1. \tag{1.9}$$

Indeed, if $\gcd\{k : r_k \neq 0\} = n > 1$, we can partition the lattice $\frac{1}{2} + \mathbb{Z}$ into n sub-lattices $\{\frac{1}{2} + i + n\mathbb{Z}\}$, $i = 0, \dots, n-1$, and separately analyze the process on each sub-lattice (since each particle cannot jump from one sub-lattice to another). In view of (1.6), throughout this paper we invoke the following scaling convention of space and time variables:

$$X = \varepsilon x, \ Y = \varepsilon y, \ x, y \in \mathbb{Z}, \ X, Y \in \varepsilon\mathbb{Z}, \quad T = \varepsilon^2 t, \ S = \varepsilon^2 s.$$

Our main result shows the convergence of Z^ε to the SHE (1.3) for any exclusion process with jump rates satisfying (1.7) and the minor assumption (1.9).

Theorem 1.1. *Equip the space $D([0,\infty), C(\mathbb{R}))$ with the topology of uniform convergence on compact intervals. Fix any $u \in (0, \frac{1}{2})$, suppose that the initial condition satisfies*

$$\|Z_0^\varepsilon(X)\|_l := \Big[\mathbb{E}\Big(|Z_0^\varepsilon(X)|^l\Big)\Big]^{1/l} \leq C e^{A_0|X|}, \qquad \text{for } 1 \leq l \leq 6, \tag{1.10a}$$

$$\|Z_0^\varepsilon(X) - Z_0^\varepsilon(Y)\|_l \leq C|X-Y|^u e^{A_0(|X|+|Y|)}, \qquad \text{for } 1 \leq l \leq 6, \tag{1.10b}$$

$$Z_0^\varepsilon(X) \text{ convereges weakly to } \mathcal{Z}_0(X), \tag{1.10c}$$

where A_0 and C are positive constants independent of ε, X, Y, and $\mathcal{Z}_0(\,\cdot\,)$ is a random field over $\mathbb{R}$ with continuous sample paths. Then the law of $Z^\varepsilon_\cdot(\,\cdot\,)$ converges weakly to a law on $C([0,\infty), C(\mathbb{R}))$. Moreover, after a time rescaling $T' = \alpha^{-1}T$, the law coincides with the mild solution to the SHE (1.3) *with initial condition $\mathcal{Z}_0$:*

$$\mathcal{Z}_{\alpha^{-1}T}(X) = \int_{\mathbb{R}} P_T(X-Y)\mathcal{Z}_0(Y)dY + \int_0^T \int_{\mathbb{R}} P_{T-S}(X-Y)\mathcal{Z}_{\alpha^{-1}S}(Y)W(dY, dS), \tag{1.11}$$

where $P_T(X) := (2\pi T)^{-1/2}\exp(-X^2/2T)$ is the heat kernel.

Remark 1.2. For the existence, uniqueness, and positivity of solutions to (1.11), see [6, Proposition 2.5] and [2, 3, 16, 24]. When stating Theorem 1.1 we implicitly extended processes defined on $\varepsilon\mathbb{Z}$ to $\mathbb{R}$ by linear interpolation. We will use this convention throughout this paper.

Utilizing the Gätner transformation, Bertini and Giacomin [3] proved Theorem 1.1 for the special case of simple exclusion process. In contrast with simple exclusion process, non-simple exclusion processes are not (yet) explicitly solvable. Nevertheless, by Theorem 1.1 and the uniqueness of SHE (1.11), the class of exclusion processes we consider all share the same scaling exponents and limit distributions. That is, we prove *universality*, which is the first result of this kind in the context of KPZ universality class.

While there is no known exact formula for correlation functions or moments for non-simple exclusion processes, we deduce exact limit statistics for two special initial conditions, by adopting Theorem 1.1 in a manner similar to that of [1, 7].

Theorem 1.3. *Consider, in terms of occupation variable η, the step initial condition*

$$\eta_0(x) = \mathbb{1}_{[0,\infty)}(x). \tag{1.12}$$

Let $\mathcal{H}^\varepsilon_T$ be the scaled height fluctuation function defined as in (1.5). *For each fixed $(X,T) \in \mathbb{R} \times (0,\infty)$, $\mathcal{H}^\varepsilon_T(X)$ converges in distribution to $\mathcal{H}_T(X)$, and $\mathcal{F}(T,X) := -(\mathcal{H}_T)^\alpha(X) + \frac{X^2}{2T} + \frac{1}{2}\log(2\pi T)$ has the same law as in* [1, Theorem 1.1]. *In particular,*

$$-\left[(\mathcal{H}_T)^\alpha(X) - \frac{X^2}{2T} - \frac{T}{24}\right](T/2)^{-\frac{1}{3}}$$

converges in distribution as $T \to \infty$ to the GUE Tracy-Widom distribution.

Theorem 1.4. *Consider, in terms of occupation variable η, the step Bernoulli initial condition*

$$\eta_0(x) = \mathbb{1}_{[0,\infty)}(x) \sum_{0 \le y \le x} (2b_y - 1), \tag{1.13}$$

where b_y, $y = 0, 1, \ldots$, are i.i.d. Bernoulli random variable with $\mathbb{P}(b_y = 1) = \frac{1}{2}$. As in [7], *define*

$$h^{fluc}_t(x) := \frac{\left(h_{\gamma^{-1}t}\right)^\alpha(x) - \frac{t}{2}}{t^{1/3}}, \; T := \varepsilon^{\frac{3}{2}} t, \; X := 2^{-\frac{1}{3}} t^{-\frac{2}{3}} x,$$

where γ is defined as in (1.6a). *h^{fluc} has the same limit law as in* [7, Theorem 3].

Remark 1.5. Adopting the proof Theorem 1.4, one may generalize this result to the rarefaction fan initial condition described in [7] and get a convergence theorem similar to [7, Theorem 16].

Remark 1.6. The limit statistics of the SHE for the step and step Bernoulli initial conditions has been proven by Amir-Corwin-Quastel [1] and Corwin-Quastel [7] respectively, based on previous work of Tracy and Widom [21, 22] (see also [19, 20, 23]). Our exact statistic results Theorem 1.3 and 1.4 then follow from the uniqueness of the SHE and [1, 7]. Note that in the context KPZ universality class Theorem 1.3 and Theorem 1.4 are the first exact limit statistics results derived out of the limit stochastic partial differential equation, and not by explicitly solving the model in question.

Remark 1.7. It not clear if in the absence of (1.7) the fluctuation of h converges as $\varepsilon \to 0$ under scaling similar to (1.5), and, if so, is it to the same limit object. It is interesting to know whether exclusions processes satisfying (1.7) are actually explicitly solvable. Such explicit solvability could provide more information, such as non-asymptotic properties, for the processes in question.

In deriving Theorem 1.1, the transformed equation does not take the same form as in the simple case. Aside from the Laplacian-type term (2.14), here a quasi-linear (2.15) term appears. This quasi-linear term carries negligible factors (see (2.12)). Hence, we apply a discrete analog of parametrix method [9, Chapter 1] in parabolic partial differential equations to control the contribution of this quasi-linear term to the solution semi-group.

Once this is done, the convergence to the SHE is reduced to showing the convergence of the quadratic variation of the corresponding martingale term (see (2.5)). This is done by analysis the integral equation (5.16), which can be view as a linear equation

$$g^\varepsilon = \mathbf{q} g^\varepsilon + L + O(\varepsilon^u)$$

in a suitable normed space, where $u \in (0, \frac{1}{2})$. We then use the key identity $L = 0$ (see (5.5)), which generalize [3, Lemma A.1], to show that

$$(I - \mathbf{q})g^\varepsilon = O(\varepsilon^u).$$

Employing the inequality [3, Lemma A.2], Bertini-Giacomin showed that the linear operator $\mathbf{q}$ is a contraction map, which implies that g^ε is small. A technical challenge appears when we extend this to $m > 1$: the operator $\mathbf{q}$ is no longer a contraction map when $m > 1$ (See Remark 5.5 and Proposition A.3.) We meet this challenge by first observing that for weak convergence we do not require the full strength of the convergence proven by [3]. Instead, we need only to show that certain space-time average of g^ε converges to zero. (See Remark 5.8.) To this end, we first reduce the original integral equation into a *finite dimensional* linear equation. For this finite dimensional equation we have a generalization (see (5.6)) of [3, Lemma A.2], which shows contractivity.

We next outline this paper. In Sections 2–5, we prove the convergence toward the SHE, namely Theorem 1.1. This is a direct consequence of the follow two propositions, which we establish in Sections 3 and 4 respectively.

Proposition 1.8. *Assume that the initial condition satisfies* (1.10a) *and* (1.10b)*, then the laws of* $\{Z^\varepsilon\}_\varepsilon$ *on* $D([0,\infty), C(\mathbb{R}))$ *is tight. Moreover, limit points of the law of* $\{Z^\varepsilon\}_\varepsilon$ *concentrate on* $C([0,\infty), C(\mathbb{R}))$*.*

Proposition 1.9. *The law of any limit point* $\mathcal{Z}$ *of* $\{Z^\varepsilon\}$ *is a mild solution* (1.11) *to the SHE.*

In Section 2, we introduce the Gärtner transformation. In Section 3, we show that the quasi-linear term have negligible effect and prove Proposition 1.8 . In Section 4, we reduce Proposition 1.9 to the convergence of the quadratic variation using the martingale problem method. In Section 5, we establish the convergence of the quadratic variation, deriving the integral equation (5.16) in Section 5.1 and reducing the integral equation to a finite dimensional linear equation in Section 5.2. In section 6, we establish initial-layer estimates for the two special initial conditions (1.12) and (1.13). The initial-layer estimates allow us to apply Theorem 1.1 to these initial conditions, and therefore get Theorem 1.3 and 1.4.

Acknowledgment. We thank Yao-Yuan Mao for the numerical prediction of Lemma 5.1, and thank Ivan Corwin and Stefano Olla for helpful conversations.

2. The Gärtner transformation.

We consider finite range non-simple exclusion process, and the corresponding growth model whose height function is given by (1.4). Let

$$\left\{P^k_T(X), Q^k_T(X) : k = 1, \ldots, m,\ X \in \varepsilon\mathbb{Z}\right\} \tag{2.1}$$

be mutually independent (in k and X) Poisson processes in T, with each $P^k_T(X)$ having rate $p_k\gamma^{-1}\varepsilon^{-3/2}$, each $Q^k_T(X)$ having rate $q_k\gamma^{-1}\varepsilon^{-3/2}$. Each $P^k_T(X)$ counts the total numbers of *attempted* right-going jumps with size k starting at $X - \frac{\varepsilon}{2}$, within the time interval $[0, T]$; Each $Q^k_T(X)$ does the same for left-going jumps.

We start to derive the governing stochastic differential equation of Z^ε. To simplify notation we omit the dependence on ε when this causes no confusion. For example we sometimes write λ_ε as λ. From the dynamics of h described in the introduction, we know that the height at x increases by two (a box) when a particle jump across x from right to left. By the definition (1.5) of Z^ε, during $[T, T + dT]$ the total contribution to $dZ^\varepsilon_T(X)$ of left-going jumps reads

$$\left(e^{-2\lambda} - 1\right) Z^\varepsilon_T(X) \sum_{k=1}^{m} \sum_{j=1}^{k} v^{jk}_t(x) dQ^k_T(X + j\varepsilon),$$

where

$$v_t^{jk}(x) := \frac{1-\eta_t(x+j-k)}{2}\,\frac{1+\eta_t(x+j)}{2} \in \{0,1\}, \tag{2.2}$$

indicating by 1 allowed jumps from site $x+j$ to site $x+j-k$, by 0 prohibited jumps from site $x+j$ to site $x+j-k$.

Similarly, the contribution of right-going jumps to $Z_T^\varepsilon(X)$ reads

$$\big(e^{2\lambda}-1\big) Z_T^\varepsilon(X) \sum_{k=1}^{m}\sum_{j=1}^{k} u_t^{jk}(x) dP_T^k(X+(j-k)\varepsilon),$$

where

$$u_t^{jk}(x) := \frac{1+\eta_t(x+j-k)}{2}\,\frac{1-\eta_t(x+j)}{2} \in \{0,1\}. \tag{2.3}$$

Aside from the contributions of jumps, there is a continuous growth coming from the term $\exp(\frac{\nu T}{\gamma\varepsilon^{3/2}})$. Gathering the preceding contributions together, and separating drifts from each P^k and Q^k, we obtain the following infinite ($X \in \varepsilon\mathbb{Z}$) system of stochastic differential equations in T:

$$dZ^\varepsilon = \Omega^\varepsilon Z^\varepsilon dT + Z^\varepsilon dM^\varepsilon, \tag{2.4}$$

where

$$\Omega_T^\varepsilon(X) = \frac{1}{\gamma\varepsilon^{\frac{3}{2}}}\left\{\nu + \sum_{k=1}^{m}\sum_{j=1}^{k}\Big[\big(e^{2\lambda}-1\big)p_k u_t^{jk}(x) + \big(e^{-2\lambda}-1\big)q_k v_t^{jk}(x)\Big]\right\}.$$

Here $M_T^\varepsilon(X)$ is a martingale in T for all $X \in \varepsilon\mathbb{Z}$, which reads

$$\begin{aligned} dM_T^\varepsilon(X) = {} & \big(e^{2\lambda}-1\big)\sum_{k=1}^{m}\sum_{j=1}^{k} u_t^{jk}(x)\Big(dP_T^k(X+(j-k)\varepsilon) - \frac{p_k}{\gamma\varepsilon^{3/2}}dT\Big) \\ & + \big(e^{-2\lambda}-1\big)\sum_{k=1}^{m}\sum_{j=1}^{k} v_t^{jk}(x)\Big(dQ_T^k(X+j\varepsilon) - \frac{q_k}{\gamma\varepsilon^{3/2}}dT\Big). \end{aligned} \tag{2.5}$$

For simple exclusion process, Gärtner [10] showed that some appropriate choice of γ, λ, ν turns the nonlinear drift term $\Omega^\varepsilon Z^\varepsilon$ in (2.4) into a discrete Laplacian. That is, $\Omega^\varepsilon Z^\varepsilon = \frac{1}{2}\Delta Z^\varepsilon$, which in particular is linear. We next verify that this property extends with modification to non-simple exclusion processes with constraint (1.7).

Using the definitions (2.3) and (2.2) of u^{jk} and v^{jk}, we have

$$\Omega^\varepsilon = \Big(\gamma\varepsilon^{3/2}\Big)^{-1}(\nu + a + b), \tag{2.6}$$

where

$$a := -\frac{1}{4}\sum_{k=1}^{m}\sum_{j=1}^{k}\big[\big(e^{2\lambda}-1\big)p_k + \big(1-e^{-2\lambda}\big)q_k\big]\Big(\eta_t(x+j) - \eta_t(x+j-k)\Big)$$

is a term linear in η, and

$$b := \frac{1}{4}\sum_{k=1}^{m}\sum_{j=1}^{k}\big[\big(e^{2\lambda}-1\big)p_k - \big(1-e^{-2\lambda}\big)q_k\big]\Big(1 - \eta_t(x+j)\eta_t(x+j-k)\Big)$$

is the major cause of non-linearity. By (1.7) and our choice (1.6b) of λ,

$$\big(e^{2\lambda}-1\big)p_k - \big(1-e^{-2\lambda}\big)q_k = 0,$$

and therefore $b = 0$. Next, we proceed to show that aZ_ε is the sum of a discrete Laplacian of Z^ε, a constant drift term, and a presumably small quasi-linear term. Indeed, under (1.7) and (1.6), a becomes

$$-\frac{\gamma}{2}\sum_{k=1}^{m} r_k \sum_{j=1}^{k}\Big(\eta_t(x+j)-\eta_t(x-j+1)\Big),$$

where we have rearranged the summation over j. Using the telescoping identity

$$\eta_t(x+j)-\eta_t(x-j+1)=\sum_{|l|<j}(\eta_t(x+l+1)-\eta_t(x+l)),$$

we have

$$a=-\frac{\gamma}{2}\sum_{k=1}^{m} r_k \sum_{|j|<k}(k-|j|)\Big(\eta_t(x+j+1)-\eta_t(x+j)\Big). \tag{2.7}$$

Define the k-steps backward discrete gradient and k-steps discrete Laplacian as

$$\begin{aligned} \nabla_k^- f(x) &:= f(x)-f(x-k), & \nabla_k^- F(X) &:= F(X)-F(X-k\varepsilon),\\ \Delta_k f(x) &:= f(x+k)+(x-k)-2f(x), & \Delta_k F(X) &:= F(X+k\varepsilon)+F(X-k\varepsilon)-2F(X). \end{aligned}$$

One-step gradient ∇_1^- and one-step Laplacian Δ_1 are simply denoted by ∇^- and Δ, respectively. Set $\nu' := e^{-\lambda}+e^{\lambda}-2$, $\nu'' := \frac{1}{2}\big(e^{\lambda}-e^{-\lambda}\big)$. From (1.5) and (1.6), we have the following identity,

$$(\Delta Z_T^\varepsilon)(X+j\varepsilon)=\Big[\nu'-\nu''\Big(\eta_t(x+j+1)-\eta_t(x+j)\Big)\Big]Z_T^\varepsilon(X+j\varepsilon), \tag{2.8}$$

which can be directly verified per possible value $(\pm1,\pm1)$ that $(\eta_t(x+j+1),\eta_t(x+j))$ takes. Multiplying both sides of (2.7) by $Z_T^\varepsilon(X)$, and combining (2.7) and (2.8) using the easily verified identities $\gamma/(2\nu'')=\frac{1}{2}(1-\varepsilon)^{1/2}$ and $\gamma\nu'/(2\nu'')=1-(1-\varepsilon)^{1/2}$, we get

$$aZ_T^\varepsilon(X)=\frac{(1-\varepsilon)^{\frac12}}{2}\sum_{k=1}^{m} r_k\sum_{|j|<k}(k-|j|)\Delta Z_T^\varepsilon(X+\varepsilon j)\frac{Z_T^\varepsilon(X)}{Z_T^\varepsilon(X+\varepsilon j)}-\alpha\Big[1-(1-\varepsilon)^{\frac12}\Big]Z_T^\varepsilon(X).$$

Next, writing $Z_T^\varepsilon(X)/Z_T^\varepsilon(\varepsilon+j\varepsilon)$ as the sum of 1 and $Z_T^\varepsilon(X)/Z_T^\varepsilon(X+j\varepsilon)-1$, using the telescoping identity

$$\sum_{|j|<k}(k-|j|)\Delta f(X+\varepsilon j)=\Delta_k f(X), \tag{2.9}$$

we compute aZ_T^ε as

$$aZ_T^\varepsilon=\frac{(1-\varepsilon)^{\frac12}}{2}\sum_{k=1}^{m} r_k\Delta_k Z_T^\varepsilon+\mathscr{E}_T^\varepsilon-\alpha\Big[1-(1-\varepsilon)^{\frac12}\Big]Z_T^\varepsilon,$$

where $\mathscr{E}^\varepsilon$ is a quasi-linear term defined as

$$\mathscr{E}_T^\varepsilon(X):=\sum_{|j|<m} b_t^{\varepsilon,j}(x)(\Delta Z_T^\varepsilon)(X+j\varepsilon),\quad b_t^{\varepsilon,j}(x):=\sum_{k>|j|} r_k\bigg(\frac{Z_T^\varepsilon(X)}{Z_T^\varepsilon(X+j\varepsilon)}-1\bigg). \tag{2.10}$$

The choice (1.6c) of ν eliminates the constant drift term $\alpha[1-(1-\varepsilon)^{1/2}]$ in (2.6). Since $|\lambda_\varepsilon|\le C\varepsilon^{1/2}$, see (1.6b), we have the following a priori bounds

$$Z_T^\varepsilon(Y)=\Big(1+O(\varepsilon^{\frac12}|x-y|)\Big)Z_T^\varepsilon(X), \tag{2.11}$$

$$|b_t^{\varepsilon,j}(x)|\le C\varepsilon^{\frac12}, \tag{2.12}$$

where C is a constant independent of ε, X, and T, and $O(\zeta)$ represents a generic function such that $\sup_{(0,\infty)}|O(\zeta)||\zeta|^{-1}\le C$. Therefore, intuitively, as long as ΔZ^ε is well-behaved we expect the contribution of $\mathscr{E}^\varepsilon$ to be negligible in the continuum limit $\varepsilon\to 0$.

Putting together the preceding calculation, we have

$$dZ^\varepsilon = \frac{1}{2\varepsilon^2}(\mathcal{L} + \mathcal{T}^\varepsilon_T) Z^\varepsilon dT + Z^\varepsilon dM^\varepsilon, \tag{2.13}$$

where the operators $\mathcal{L}$, $\mathcal{T}^\varepsilon_T$ act on Z^ε_T as

$$\Big(\mathcal{L} Z^\varepsilon_T\Big)(X) = (1-\varepsilon)^{\frac{1}{2}} \sum_{k=1}^{m} r_k (\Delta_k Z^\varepsilon_T)(X), \tag{2.14}$$

$$\Big(\mathcal{T}^\varepsilon_T Z^\varepsilon_T\Big)(X) = (1-\varepsilon)^{\frac{1}{2}} \sum_{|j|<m} b^{\varepsilon,j}_t(x)(\Delta Z^\varepsilon_T)(X + j\varepsilon). \tag{2.15}$$

3. ESTIMATES OF THE DRIFT TERM.

In this section we focus on the drift term $(2\varepsilon^2)^{-1}(\mathcal{L} + \mathcal{T}^\varepsilon_T)Z^\varepsilon$ of (2.13). Since $\varepsilon \to 0$, we remove the $(1-\varepsilon)^{1/2}$ factor (2.14) and (2.15) by a time rescaling $T' = (1-\varepsilon)^{1/2}T$, without affecting the limit distribution. Since we only need weak convergence, throughout this paper we work in a bounded macroscopic time interval $T \in [0, \tilde{T}]$, with $\tilde{T}$ arbitrary but fixed. Throughout our analysis C refers to a finite positive constant. It may depend on various parameters (i.e. $\tilde{T}$ or the bounds (1.10)), but is always independent of ε, X, and $T \in [0, \tilde{T}]$. We use $*$ to denote macroscopic convolution and $\star$ to denote microscopic convolution, that is

$$(F * G)(X) := \varepsilon \sum_{Y \in \varepsilon\mathbb{Z}} F(X-Y)G(Y), \quad (f \star g)(x) := \sum_{y\in\mathbb{Z}} f(x-y)g(y).$$

Let $\mathbf{p}$, $\mathbf{P}^\varepsilon$ be the microscopic and macroscopic semigroups associated with $\mathcal{L}$, respectively:

$$\frac{d}{dt}\mathbf{p}_t(x) = \frac{1}{2}\mathcal{L}\mathbf{p}_t(x), \quad \mathbf{p}_0(x) = \mathbb{1}_{\{0\}}(x), \tag{3.1}$$

$$\mathbf{P}^\varepsilon_T(X) = \frac{1}{\varepsilon}\mathbf{p}_{\varepsilon^{-2}T}\big(\varepsilon^{-1}X\big). \tag{3.2}$$

Similarly, define $\tilde{\mathbf{P}}^\varepsilon$ as the fundamental solution of the following discrete partial differential equation:

$$\frac{d}{dT}\tilde{\mathbf{P}}^\varepsilon_{T,S}(X,Y) = \frac{1}{2\varepsilon^2}(\mathcal{L} + \mathcal{T}^\varepsilon_T)\tilde{\mathbf{P}}^\varepsilon_{T,S}(X,Y), \quad \tilde{\mathbf{P}}^\varepsilon_{S,S}(X,Y) = \frac{1}{\varepsilon}\mathbb{1}_{\{Y\}}(X), \tag{3.3}$$

where the operators $\mathcal{L}$, $\mathcal{T}^\varepsilon_T$ act on the first space variable. Note that $\tilde{\mathbf{P}}^\varepsilon$ is random because $b^{\varepsilon,j}$ is, whereas $\mathbf{P}^\varepsilon$ is deterministic. The discrete SHE (2.13) can be expressed in integrated form as

$$Z^\varepsilon_T = \tilde{\mathbf{P}}^\varepsilon_{T,0} * Z^\varepsilon_0 + \int_0^T \tilde{\mathbf{P}}^\varepsilon_{T,S} * (Z^\varepsilon_S dM^\varepsilon_S). \tag{3.4}$$

When $m = 1$, since j takes only the value 0, for which $b^{\varepsilon,0}_t = 0$, $\tilde{\mathbf{P}}^\varepsilon$ equals the discrete heat kernel $\mathbf{P}^\varepsilon$, which can be analyzed directly. In general,

$$\widehat{\mathbf{P}}^\varepsilon_{T,S}(X,Y) := \tilde{\mathbf{P}}^\varepsilon_{T,S}(X,Y) - \mathbf{P}^\varepsilon_{T-S}(X-Y) \neq 0.$$

Using the a priori bound (2.12), we show in the sequel that $\widehat{\mathbf{P}}^\varepsilon$ to decay to zero as $\varepsilon \to 0$ by estimating $\widehat{\mathbf{P}}^\varepsilon$ directly.

Given any $A \geq 0$ and $n \in \mathbb{N}$, define the norms

$$|F|_A := \sup_X e^{-A|X|}\varepsilon \sum_Y \|F(X,Y)\|_\infty e^{A|Y|}, \quad [G]_{A,n} := \sup_X \|G(X)\|_n e^{-A|X|}, \tag{3.5}$$

for any random field $F(X,Y)$ and $G(X)$, noting that for any random field $F'(X,Y)$

$$|F * F'|_A \leq |F|_A |F'|_A, \tag{3.6}$$

$$\sup_{X,Y} \|(F * F')(X,Y)\|_\infty \leq \Big(\sup_{X,Y}\|F(X,Y)\|_\infty\Big)|F'|_0, \tag{3.7}$$

$$e^{-|A|X}\varepsilon\sum_{Y}\|F(X,Y)\|_{\infty}\|G(Y)\|_{n}\leq|F|_{A}[G]_{A,n}. \tag{3.8}$$

Proposition 3.1. *Given any $A\geq 0$, $u\in(0,\frac{1}{2})$, $v\in[0,1]$ there exists a constant C such that*

$$|\widehat{\mathbf{P}}^{\varepsilon}_{T,S}|_{A}\leq\varepsilon^{u}C, \tag{3.9}$$

$$\left\|\widehat{\mathbf{P}}^{\varepsilon}_{T,S}(X,Y)\right\|_{\infty}\leq(T-S)^{-\frac{1}{2}}\varepsilon^{u}C, \tag{3.10}$$

$$\left\|\widehat{\mathbf{P}}^{\varepsilon}_{T,S}(X,Y)-\widehat{\mathbf{P}}^{\varepsilon}_{T,S}(X',Y)\right\|_{\infty}\leq\left(|X-X'|(T-S)^{-\frac{1}{2}}\right)^{v}(T-S)^{-\frac{1}{2}}\varepsilon^{u}C, \tag{3.11}$$

whenever ε is small enough, where $0\leq S\leq T\leq\tilde{T}$.

Proof. Let $\mathbf{D}_{T,S}(X,Y):=(2\varepsilon^{2})^{-1}\big(\mathcal{T}^{\varepsilon}_{T}\mathbf{P}^{\varepsilon}_{T-S}\big)(X,Y)$, and

$$\Delta^{T}_{S}(n):=\{(T_1,\ldots,T_n)\in\mathbb{R}^{n}:S:=T_{n+1}<T_n<\ldots<T_1<T\}. \tag{3.12}$$

The equation (3.3) is equivalent to

$$\tilde{\mathbf{P}}^{\varepsilon}_{T,S}=\mathbf{P}^{\varepsilon}_{T-S}+\int_{S}^{T}\mathbf{P}^{\varepsilon}_{T-T'}*(2\varepsilon^{2})^{-1}\mathcal{T}^{\varepsilon}_{T'}\tilde{\mathbf{P}}^{\varepsilon}_{T',S}dT'. \tag{3.13}$$

Let

$$\mathbf{B}^{n}_{T,S}:=\int_{\Delta^{T}_{S}(n)}\mathbf{P}^{\varepsilon}_{T-T_1}*\mathbf{D}_{T_1,T_2}*\cdots*\mathbf{D}_{T_n,S}\,dT_1\cdots dT_n.$$

Iterating (3.13), we obtain $\widehat{\mathbf{P}}^{\varepsilon}=\sum_{n=1}^{\infty}\mathbf{B}^{n}$ (which is well-defined since as we show in the sequel the infinite sum converges in the norm $|\cdot|_{A}$). From this expression of $\mathbf{B}^{n}$, using (3.6) and (3.7) we have

$$\left|\mathbf{B}^{n}_{T,S}\right|_{A}\leq\int_{S}^{T}\left|\mathbf{P}^{\varepsilon}_{T-T_1}\right|_{A}D^{n}_{T_1,S}(A)\,dT_1, \tag{3.14a}$$

$$\sup_{X,Y}\left\|\mathbf{B}^{n}_{T,S}(X,Y)\right\|_{\infty}\leq\int_{S}^{T}\Big(\sup_{X}\mathbf{P}^{\varepsilon}_{T-T_1}(X)\Big)D^{n}_{T_1,S}(0)\,dT_1, \tag{3.14b}$$

$$\sup_{Y}\left\|\nabla^{-}_{n}\mathbf{B}^{n}_{T,S}(X,Y)\right\|_{\infty}\leq\int_{S}^{T}\Big(\sup_{X}\left\|\nabla^{-}_{n}\mathbf{P}^{\varepsilon}_{T-T_1}(X)\right\|_{\infty}\Big)D^{n}_{T_1,S}(0)\,dT_1, \tag{3.14c}$$

where

$$D^{n}_{T_1,S}(A):=\int_{\Delta^{T_1}_{S}(n-1)}\prod_{j=1}^{n}\left|\mathbf{D}_{T_j,T_{j+1}}\right|_{A}dT_j. \tag{3.15}$$

We now proceed to estimate $D^{n}_{T,T_1}(A)$. By the definition of the norm $|\cdot|_{A}$ and (2.12)

$$\left|\mathbf{D}_{T_j,T_{j+1}}\right|_{A}\leq C\varepsilon^{\frac{1}{2}}\sup_{X}\Big(\varepsilon\sum_{Y}\Big|\varepsilon^{-2}\Delta\mathbf{P}^{\varepsilon}_{T_j-T_{j+1}}(X-Y)\Big|e^{A|X-Y|}\Big). \tag{3.16}$$

For any $\delta\in(0,1)$, the bound (A.10) yields

$$\Big|\varepsilon^{-2}\Delta\mathbf{P}^{\varepsilon}_{T_j-T_{j+1}}(X-Y)\Big|\leq\varepsilon^{-3}C(\delta)\exp\big(-(A+1+\delta^{-1})|X-Y|\big),$$

where we highlight the dependence of C on δ. Therefore, when $|X-Y|>\delta|\log\varepsilon|$,

$$\Big|\varepsilon^{-2}\Delta\mathbf{P}^{\varepsilon}_{T_j-T_{j+1}}(X-Y)\Big|\leq Ce^{-(A+1)|X-Y|}. \tag{3.17}$$

When $0<|X-Y|\leq\delta|\log\varepsilon|$, using (A.9) we obtain

$$\left|\varepsilon^{-2}\Delta\mathbf{P}^{\varepsilon}_{T}(X-Y)\right|\leq|X-Y|^{-1}\big(T\vee\varepsilon^{-2}\big)^{-1}C. \tag{3.18}$$

When $X=Y$, using (A.7) we obtain

$$\left|\varepsilon^{-2}\Delta\mathbf{P}^{\varepsilon}_{T}(X-Y)\right|\leq\big(T\vee\varepsilon^{-2}\big)^{-\frac{3}{2}}C. \tag{3.19}$$

Breaking the sum in (3.16) into $|X-Y| > \delta|\log\varepsilon|$, $0 < |X-Y| \leq \delta|\log\varepsilon|$, and $X = Y$, and using (3.17), (3.18), and (3.19) respectively, we see that the sum is dominated by contribution of the middle part, yielding

$$\big|\mathbf{D}_{T_j,T_{j+1}}\big|_A \leq |\log\varepsilon|\varepsilon^{\frac{1}{2}-A\delta}\big(T\vee\varepsilon^{-2}\big)^{-1}C(\delta).$$

Plugging this estimate back to (3.15) and evaluating the time integrals using the readily verified estimate

$$\int_0^T \big(S\vee\varepsilon^{-2}\big)^{-1}\big((T-S)\vee\varepsilon^{-2}\big)^{-1}dS \leq C|\log\varepsilon|\big(T\vee\varepsilon^{-2}\big)^{-1}, \text{ for } T\in[0,\tilde{T}],$$

we get

$$D^n_{T_1,S} \leq \Big(\varepsilon^{\frac{1}{2}-A\delta}|\log\varepsilon|^2C(\delta)\Big)^n\big(T\vee\varepsilon^{-2}\big)^{-1}. \tag{3.20}$$

The bound (A.10) implies

$$|\mathbf{P}^\varepsilon_{T'}|_A \leq C. \tag{3.21}$$

Applying the estimate (3.20) of $D^n_{T_1,S}$ and the estimates (3.21), (A.4), (A.6) of $\mathbf{P}^\varepsilon$ to (3.13), evaluation the time integral, and fixing some $\delta < (\frac{1}{2}-u)A^{-1}$, we get

$$\begin{aligned}&\big|\mathbf{B}^n_{T,S}\big|_A \leq (\varepsilon^uC)^n, \quad \big\|\mathbf{B}^n_{T,S}\big\|_\infty \leq (T-S)^{-\frac{1}{2}}(\varepsilon^uC)^n,\\ &\big\|\nabla^-_n\mathbf{B}^n_{T,S}\big\|_\infty \leq \Big(|X-X'|(T-S)^{-\frac{1}{2}}\Big)^v\big((T-S)\vee\varepsilon^{-2}\big)^{-\frac{1}{2}}(\varepsilon^uC)^n.\end{aligned}$$

Summing these geometric series proves the Proposition. □

The estimates (3.9), (3.10), and (3.11) together with (3.21), (A.4) and (A.6) directly imply the following corollary.

Corollary 3.2. *For any $A\geq 0$, $0\leq S\leq T\leq\tilde{T}$, there exists a constant C such that*

$$\sup_{0\leq S\leq T\leq\tilde{T}}\Big|\tilde{\mathbf{P}}_{T,S}\Big|_A \leq C, \tag{3.22}$$

$$\sup_{X,Y}\Big\|\tilde{\mathbf{P}}^\varepsilon_{T,S}(X,Y)\Big\|_\infty \leq (T-S)^{-\frac{1}{2}}C, \tag{3.23}$$

$$\sup_Y\Big\|\tilde{\mathbf{P}}^\varepsilon_{T,S}(X,Y)-\tilde{\mathbf{P}}^\varepsilon_{T,S}(X',Y)\Big\|_\infty \leq \Big(|X-X'|(T-S)^{-\frac{1}{2}}\Big)^v(T-S)^{-\frac{1}{2}}C, \tag{3.24}$$

whenever ε is small enough.

We proceed to establish Hölder continuities of Z^ε in space and time. They lead directly to tightness of Z^ε as stated in Proposition 1.8. Moreover, they are used in establishing the convergence of the martingale term, in the course of proving Proposition 5.4. First we need a technical lemma.

Lemma 3.3. *Given any random field $F_S(X,Y)$, let $R_{T,T'} := \int_{T'}^T F_S * (Z^\varepsilon_S dM^\varepsilon_S)$ For any $A\geq 0$, $n\in\mathbb{N}$ we have*

$$\big\|R_{T,T'}(X)^2\big\|_n \leq C\int_{T'}^T \varepsilon\sum_Y\big(\mathcal{F}(F)\big)_S(X,Y)\,\big\|Z^\varepsilon_S(Y)^2\big\|_n dS,$$

where

$$\big(\mathcal{F}(F)\big)_S(X,Y) := \sup_{S'\in\mathcal{I}_S}\sup_{|l|<m}\big\|F_{S'}(X,Y+l\varepsilon)^2\big\|_\infty,$$

$$\mathcal{I}_S := \Big(\varepsilon^2\lceil\varepsilon^{-2}S\rceil-1, \varepsilon^2\lceil\varepsilon^{-2}S\rceil\Big] \subset \mathbb{R}.$$

Proof. By the BDG inequality,

$$\big\| R_{T,T'}(X)^2 \big\|_n \leq C \big\| [R_{\cdot,T'}(X), R_{\cdot,T'}(X)]_T \big\|_n. \tag{3.25}$$

Let $\mathfrak{T}_{(T',T]}(y)$ be the *random* set of all $S \in (T', T]$ at which a jump across the site $y - \frac{1}{2}$ occurs. Since the Poission processes defined in (2.1) are independent, using (2.5) we have

$$\begin{aligned} &[R_{\cdot,T'}(X), R_{\cdot,T'}(X)]_T \\ &= \varepsilon^2 \sum_{Y} \sum_{|l|<m} \sum_{S \in \mathfrak{T}(y,l)} F'_S(X,Y) F'_S(X, Y + l\varepsilon)(e^{2\sigma(y,l)\lambda} - 1)^2 Z^\varepsilon_{T'}(Y) Z^\varepsilon_{T'}(Y + l\varepsilon), \end{aligned}$$

where $\mathfrak{T}^l(y) := \mathfrak{T}_{(T',T]}(y) \cap \mathfrak{T}_{(T',T]}(y + l)$ is the set of $S \in (T', T]$ at which a jump across *both* the site $y - \frac{1}{2}$ and $y + l - \frac{1}{2}$ occurs, and $\sigma(y, l) = 1$ when the jump goes the right, $\sigma(y, l) = -1$ when the jump goes the left. Note that we need only to sum over $|l| < m$ since $\mathfrak{T}^l(y) = \emptyset$ when $|l| \geq m$. Partition $(0, \infty)$ into subintervals $\mathcal{I}_i := (\varepsilon^{-2} i, \varepsilon^{-2}(i + 1)]$. Using $|e^{\pm\lambda} - 1| \leq C\varepsilon^{1/2}$, and replacing F_S and Z^ε_S by their supremum over $\mathcal{I}_i$, we have

$$[R_{\cdot,T'}(X), R_{\cdot,T'}(X)]_T \leq C\varepsilon^3 \sum_{i=\lfloor t' \rfloor}^{\lfloor t \rfloor} \sum_{Y} \mathcal{F}(F')_{i\varepsilon^2}(X,Y) \max_{|l|<m} \sup_{S \in \mathcal{I}_i} Z^\varepsilon_S(Y + l\varepsilon)^2. \tag{3.26}$$

Since each jump increase or decrease Z^ε_T by a factor of $e^{2\lambda}$,

$$\sup_{S \in \mathcal{I}_i} Z^\varepsilon_S(Y)^2 \leq e^{4\lambda Q_i(y)} Z^\varepsilon_{\varepsilon^2 i}(Y)^2, \quad \inf_{S \in \mathcal{I}_i} Z^\varepsilon_S(Y)^2 \geq e^{-4\lambda Q_i(y)} Z^\varepsilon_{\varepsilon^2 i}(Y)^2,$$

where

$$Q_i(y) := \#\mathfrak{T}_{\mathcal{I}_i}(y), \tag{3.27}$$

which is stochastically bounded by a Poisson random variable with parameter less than or equal to $\sum_{k=1}^m k r_k$. Since $Q_i(y)$ and $Z^\varepsilon_{\varepsilon^2 i}(Y)$ are independent, $0 < \mathbb{E} e^{k Q^l_i(y)} < \infty$ for any $k \in \mathbb{N}$, and $\lambda \leq C\varepsilon^{1/2} \leq C$, from the above inequalities we obtain

$$\Big\| \sup_{S \in \mathcal{I}_i} Z^\varepsilon_S(Y)^2 \Big\|_n \leq C \Big\| \inf_{S \in \mathcal{I}_i} Z^\varepsilon_S(Y)^2 \Big\|_n \leq C\varepsilon^{-2} \int_{\mathcal{I}_i} \big\| Z^\varepsilon_{T'}(Y)^2 \big\|_n dS. \tag{3.28}$$

Combining (3.25), (3.26), and (3.28), we conclude the lemma. ☐

Lemma 3.4. *Given any $k \in \mathbb{N}$, suppose the initial condition Z^ε_0 satisfies* (1.10) *for $1 \leq l \leq 2k$. For any $u \in (0, \frac{1}{2})$,*

$$\|Z^\varepsilon_T(X)\|_{2k} \leq C e^{A_0|X|}, \tag{3.29}$$

$$\|Z^\varepsilon_T(X) - Z^\varepsilon_T(X')\|_{2k} \leq |X - X'|^u C e^{A_0(|X|+|X'|)}, \tag{3.30}$$

$$\|Z^\varepsilon_T(X) - Z^\varepsilon_{T'}(X)\|_{2k} \leq \big(|T - T'|^{\frac{u}{2}} + \varepsilon^u\big) C e^{2A_0|X|}, \tag{3.31}$$

whenever ε is small enough, where A_0 is the same as in (1.10a) *and* (1.10b).

Proof. The estimate (3.29) is equivalent to $[(Z^\varepsilon_T)^2]_{2A_0,k} \leq C$, which we next show. Using (3.4) and $(a + b)^2 \leq 2a^2 + 2b^2$ we have

$$\big[(Z^\varepsilon_T)^2\big]_{2A_0,k} \leq 2\Big[(\tilde{\mathbf{P}}^\varepsilon_{T,0} * Z^\varepsilon_0)^2\Big]_{2A_0,k} + 2\Big[\Big(\int_0^T \tilde{\mathbf{P}}^\varepsilon_{T,S} * Z^\varepsilon_S dM^\varepsilon_S\Big)^2\Big]_{2A_0,k}.$$

A direct expansion using Cauchy-Schwartz inequality shows that the first term on the RHS is less than or equal to

$$2 \sup_X e^{-2A|X|} \varepsilon^2 \sum_{Y_1,Y_2} \|\tilde{\mathbf{P}}^\varepsilon_{T,0}(X, Y_1)\|_\infty \, \|\tilde{\mathbf{P}}^\varepsilon_{T,0}(X, Y_2)\|_\infty \, \|Z^\varepsilon_0(Y_1)\|_{2k} \, \|Z^\varepsilon_0(Y_2)\|_{2k},$$

which by (1.10a) and (3.22) is bounded. Now apply Lemma 3.3 for $F_S = \tilde{\mathbf{P}}^\varepsilon_{T,S}$ to estimate the second term on the RHS. Using (3.22) and (3.23) we have

$$\Big|\mathcal{F}(\tilde{\mathbf{P}}^\varepsilon_{T,\cdot})_S\Big|_{2A_0} \leq (T-S)^{-\frac{1}{2}}C, \tag{3.32}$$

which together with (3.8) implies

$$\Big[\Big(\int_0^T \tilde{\mathbf{P}}^\varepsilon_{T,S} * Z^\varepsilon_S dM^\varepsilon_S\Big)^2\Big]_{2A_0,k} \leq C\int_0^T (T-S)^{-\frac{1}{2}}\big[(Z^\varepsilon_S)^2\big]_{2A_0,k}dS.$$

Combing our preceding estimates of the first and the second terms, we have

$$\big[(Z^\varepsilon_T)^2\big]_{2A_0,k} \leq C + C\int_0^T (T-S)^{-\frac{1}{2}}\big[(Z^\varepsilon_S)^2\big]_{2A_0,k}dS,$$

which after iteration implies $\big[(Z^\varepsilon_T)^2\big]_{2A_0,k} \leq C$.

Turing to (3.30), set $n := \varepsilon^{-1}(X'-X) \in \mathbb{Z}$. View $\nabla^-_n Z^\varepsilon_T(X,X') = Z^\varepsilon_T(X) - Z^\varepsilon_T(X')$ as a function of X and X', and $\nabla^-_n \tilde{\mathbf{P}}^\varepsilon_{T,S}(X,X',Y) = \tilde{\mathbf{P}}^\varepsilon_T(X,Y) - \tilde{\mathbf{P}}^\varepsilon_{T,S}(X',Y)$ as a function of X, X', and Y. Since $\nabla^-_n Z^\varepsilon_T = 0$ when $n=0$, without lost of generality we assume $n \neq 0$. By (3.4) we have have

$$\big\|\nabla^-_n Z^\varepsilon_T(X,X')\big\|_{2k} \leq \Big\|\Big(\nabla^-_n \tilde{\mathbf{P}}^\varepsilon_{T,0} * Z^\varepsilon_0\Big)(X,X')\Big\|_{2k} + \Big\|\int_0^T \Big(\nabla^-_n \tilde{\mathbf{P}}^\varepsilon_{T,S} * (Z^\varepsilon_S dM^\varepsilon_S)\Big)(X,X')\Big\|_{2k}.$$

Let I_1 and I_2 denote the first and second terms on the RHS, respectively. For I_1 we first have

$$\Big|\nabla^-_n \tilde{\mathbf{P}}^\varepsilon_{T,0} * Z^\varepsilon_0\Big| \leq \Big|\widehat{\mathbf{P}}^\varepsilon_{T,0} * Z^\varepsilon_0\Big| + \Big|(\widehat{\mathbf{P}}^\varepsilon_{T,0})^n * Z^\varepsilon_0\Big| + \big|\nabla^-_n \mathbf{P}^\varepsilon_T * Z^\varepsilon_0\big|,$$

where $(\widehat{\mathbf{P}}^\varepsilon_{T,S})^n(X,Y) := \widehat{\mathbf{P}}^\varepsilon_{T,S}(X+n\varepsilon,Y)$. Using (3.9) and (1.10a) we get $\|\widehat{\mathbf{P}}^\varepsilon_{T,0} * Z^\varepsilon_0(X)\|_{2k} \leq e^{A_0|X|}\varepsilon^u C$, and $\|\widehat{\mathbf{P}}^\varepsilon_{T,0} * Z^\varepsilon_0(X')\|_{2k} \leq \varepsilon^u e^{A_0|X'|}C$. Using discrete integration by parts, (3.21), and (1.10b) we obtain

$$\big\|\nabla^-_n \mathbf{P}^\varepsilon_T * Z^\varepsilon_0(X,X')\big\|_{2k} = \big\|\mathbf{P}^\varepsilon_T * \nabla^-_n Z^\varepsilon_0(X,X')\big\|_{2k} \leq |X-X'|^u e^{A_0(|X|+|X'|)}C. \tag{3.33}$$

Hence $I_1 \leq (\varepsilon^u + |X-X'|^u)e^{A_0(|X|+|X'|)}C$. Next, we apply Lemma 3.3 for $F_S = \nabla^-_n \tilde{\mathbf{P}}^\varepsilon_{T,S}$ to estimate I_2. Using (3.24), (3.22), (3.29) (which we just proved), and (3.8), we conclude

$$I_2 \leq (\varepsilon^u + |X-X'|^u)e^{A_0(|X|+|X'|)}C.$$

Since $n \neq 0$, (3.30) follows.

Turning to (3.31), assuming without lost of generality $T' < T$, from (3.4) we similarly have

$$[Z^\varepsilon_T - Z^\varepsilon_{T'}]_{2A_0,2k} \leq [J_1]_{2A_0,2k} + [J_2]_{2A_0,2k} + [J_3]_{2A_0,2k},$$

where

$$J_1 := (\tilde{\mathbf{P}}^\varepsilon_{T,0} - \tilde{\mathbf{P}}^\varepsilon_{T',0}) * Z^\varepsilon_0, \quad J_2 := \int_{T'}^T \tilde{\mathbf{P}}^\varepsilon_{T,S} * Z^\varepsilon_S dM^\varepsilon_S, \quad J_3 := \int_0^{T'} (\tilde{\mathbf{P}}^\varepsilon_{T,S} - \tilde{\mathbf{P}}^\varepsilon_{T',S}) * Z^\varepsilon_S dM^\varepsilon_S.$$

For J_1, writing $\tilde{\mathbf{P}}^\varepsilon$ as the sum of $\mathbf{P}^\varepsilon$ and $\widehat{\mathbf{P}}^\varepsilon$, and using the semi-group property of $\mathbf{P}^\varepsilon$, we have

$$J_1 \leq (|\widehat{\mathbf{P}}^\varepsilon_{T,0}| + |\widehat{\mathbf{P}}^\varepsilon_{T',0}|) * Z^\varepsilon_0 + \big|\mathbf{P}^\varepsilon_{T-T'} * G_{T'} - G_{T'}\big|,$$

where $G_{T'}(X) := \mathbf{P}^\varepsilon_{T'}(X)$. From (3.9), (1.10a), and (3.8) we deduce

$$\Big[(|\widehat{\mathbf{P}}^\varepsilon_{T,0}| + |\widehat{\mathbf{P}}^\varepsilon_{T',0}|) * Z^\varepsilon_0\Big]_{2A_0,2k} \leq \Big[(|\widehat{\mathbf{P}}^\varepsilon_{T,0}| + |\widehat{\mathbf{P}}^\varepsilon_{T',0}|) * Z^\varepsilon_0\Big]_{A_0,2k} \leq \varepsilon^u C.$$

Since $\varepsilon\sum_Y \mathbf{P}^\varepsilon(X-Y) = 1$, we have

$$\big(\mathbf{P}^\varepsilon_{T-T'} * G_{T'} - G_{T'}\big)(X) = \varepsilon\sum_Y \mathbf{P}^\varepsilon_{T-T'}(X-Y)(G_{T'}(Y) - G_{T'}(X)),$$

which together with (3.33) and (A.11) implies $[\mathbf{P}^{\varepsilon}_{T-T'} * G_{T'} - G_{T'}]_{2A_0,2k} \leq ((T-T') \vee \varepsilon^2)^{\frac{u}{2}} C$. Hence $[J_1]_{2A_0,2k} \leq (\varepsilon^u + (T-T')^{\frac{u}{2}})C$. For J_2, applying Lemma 3.3 for $F_S = \tilde{\mathbf{P}}^{\varepsilon}_{T,S}$ together with (3.32), (3.29), and (3.8), we get $[J_2]_{2A_0,2k} \leq (T-T')^{1/4}C$. For J_3, we apply Lemma 3.3 for $F_S = \tilde{\mathbf{P}}^{\varepsilon}_{T,S} - \tilde{\mathbf{P}}^{\varepsilon}_{T',S}$. Note that by definition $\mathcal{F}(F+F') \leq 2\mathcal{F}(F) + 2\mathcal{F}(F')$, yielding

$$\Big|\mathcal{F}\Big(\tilde{\mathbf{P}}^{\varepsilon}_{T,\cdot} - \tilde{\mathbf{P}}^{\varepsilon}_{T',\cdot}\Big)_S\Big| \leq 4\Big|\mathcal{F}\Big(\widehat{\mathbf{P}}^{\varepsilon}_{T,\cdot}\Big)_S\Big| + 4\Big|\mathcal{F}\Big(\widehat{\mathbf{P}}^{\varepsilon}_{T,\cdot}\Big)_S\Big| + 2\big|\mathcal{F}\big(\mathbf{P}^{\varepsilon}_{T-\cdot} - \mathbf{P}^{\varepsilon}_{T'-\cdot}\big)_S\big|,$$

which together with (3.10), (3.9), (A.8), (3.21), (3.29), and (3.8) implies $[J_3]_{2A_0,2k} \leq (\varepsilon^u + (T-T')^{u/2})C$. ☐

Proof of Proposition 1.8. This proposition is the generalization of the first half (tightness) of [3, Theorem 3.3] to the non-simple case $m > 1$. The original proof by [3] for the case $m = 1$ actually applies to generic processes satisfying the conclusions of [3, Lemma 4.1, 4.5-4.7], whose proofs relies only on the conclusions of [3, Lemma 4.1-4.3] and the fact that $Q_i(y)$, as defined in (3.27), is stochastically bounded by a Poisson random variable with a fixed parameter.

In our case, Z^{ε} satisfies (3.29), (3.30), and (3.31), which correspond to the conclusions of [3, Lemma 4.1, 4.2, 4.3], respectively, except that in (3.31) the additive error of order $\varepsilon^{\frac{1}{2}-b}$ replaces ε^{1-b} in [3, Equation (4.42)], for any $b > 0$. Nevertheless, the same argument leading [3, Lemma 4.1-4.3] to [3, Lemma 4.5-4.7] still applies. Also, $Q_i(y)$ is stochastically bounded by a Poisson random variable with the parameter $(\sum_{k=1}^{m} r_k k)$, as shown in the proof of Lemma 3.3.

☐

4. Martingale Problem and Convergence to the SHE.

The key to the proof of Proposition 1.9 is the following proposition, which establishes the convergence of the predictable quadratic variation of dM^{ε} to that of $\alpha^{1/2}\dot{W}$. The proof is deferred to Section 5.

Proposition 4.1. *For any compactly supported C^{∞}-function ϕ defined on $\mathbb{R}$,*

$$B^{\varepsilon}_T(\phi) := \int_0^T \varepsilon^2 \sum_{X,Y} \phi(X)\phi(Y) Z^{\varepsilon}_S(X) Z^{\varepsilon}_S(Y) d\langle M^{\varepsilon}(X), M^{\varepsilon}(Y)\rangle_S \tag{4.1}$$

converges weakly, as laws on $C[0,\infty)$, to $\alpha \int_0^T \big(\phi^2, \mathcal{Z}^2_S\big) dS$.

To prove Proposition 1.9, recall that[3, Proposition 4.11] the SHE (1.3) is equivalent to the following martingale problem.

Definition 4.2. *Let $\mathscr{P}$ be a probability measure on $C([0,\infty), C(\mathbb{R}))$ satisfying the following condition: given any $\tilde{T} > 0$, there exists $A \geq 0$ such that*

$$\sup_{T \in [0,\tilde{T}]} \sup_X e^{-A|X|} \mathbb{E}_{\mathscr{P}}\big(Z_T(X)^2\big) < \infty. \tag{4.2}$$

The measure $\mathscr{P}$ solves the martingale problem with initial condition $\mathcal{Z}_0$ if $\mathscr{P}(Z_0 \in A) = \mathbb{P}(\mathcal{Z}_0 \in A)$ for all Borel $A \subset C(\mathbb{R})$, and

$$N_T(\phi) := (\phi, Z_T) - (\phi, Z_0) - \frac{\alpha}{2}\int_0^T \big(\partial_X^2 \phi, Z_S\big) dS, \tag{4.3}$$

$$\Lambda_T(\phi) := N_T(\phi)^2 - \alpha \int_0^T (\phi^2, Z_S^2) dS, \tag{4.4}$$

are local $\mathscr{P}$-martingale for any compactly supported C^{∞}-function ϕ defined on $\mathbb{R}$.

Specifically, we provide the statement of [3, Proposition 4.11].

Proposition 4.3 (Proposition 4.11 of [3]). *For any initial condition $\mathcal{Z}_0$ satisfying* (1.10a) *(with ε removed), the preceding martingale problem has a unique solution. Moreover, the α-scaled solution coincides with the law of the solution to* (1.11) *with initial condition $\mathcal{Z}_0$.*

From now on we use $\mathscr{C}^{(k)}$ to represent a generic stochastic process over $[0,\infty)$ satisfying $\|\mathscr{C}^{(k)}_T\|_k \leq C$, or a generic random field over $[0,\infty)\times\mathbb{R}$ satisfying $\|\mathscr{C}^{(k)}_T(X)\|_k \leq Ce^{A|X|}$, for some $A\geq 0$.

Equipped with Proposition 4.3, we proceed to establish Proposition 1.9 via the martingale problem Definition 4.2.

Proof of Proposition 1.9. Let $\mathcal{Z}$ be a weak limit point of $\{Z^\varepsilon\}$. Passing to the relevant subsequence we assume $Z^\varepsilon \Rightarrow \mathcal{Z}$. We show that the law of $\mathcal{Z}$ satisfies the martingale problem Definition 4.2. Clearly, (4.2) holds because of (3.29). Regarding (4.3), we set

$$N^\varepsilon_T(\phi) := (\phi, Z^\varepsilon_T)_\varepsilon - (\phi, Z^\varepsilon_0)_\varepsilon - \frac{1}{2}\int_0^T \big(\varepsilon^{-2}\mathcal{L}\phi, Z^\varepsilon_S\big)_\varepsilon dS, \tag{4.5}$$

where $(\phi, Z^\varepsilon_T)_\varepsilon := \varepsilon\sum Z^\varepsilon_T(X)\phi(X)$. We first plug (3.4) into the first and third terms on the RHS of (4.5), and use (3.9), (1.10a), and (3.29) to replace $\tilde{\mathbf{P}}^\varepsilon$ by $\mathbf{P}^\varepsilon$, at the cost of an error of the form $\varepsilon^u\mathscr{C}^{(2)}$. Then we use discrete integration by parts to move the operator $\mathcal{L}$ from ϕ to $\mathbf{P}^\varepsilon$, and further use the defining equations (3.1) and (3.2) of $\mathbf{P}^\varepsilon$ to get $\partial_S\mathbf{P}^\varepsilon_{S-S'}$. We get

$$\begin{aligned}N^\varepsilon_T(\phi) &= (\phi, \mathbf{P}^\varepsilon_T * Z^\varepsilon_0)_\varepsilon + \Big(\phi, \int_0^T \mathbf{P}^\varepsilon_{T-S} * (Z^\varepsilon_S dM^\varepsilon_S)\Big)_\varepsilon - (\phi, Z^\varepsilon_0)_\varepsilon \\ &\quad - \int_0^T (\phi, (\partial_S\mathbf{P}^\varepsilon_S) * Z^\varepsilon_0)_\varepsilon - \int_0^T \Big(\phi, \int_0^S (\partial_S\mathbf{P}^\varepsilon_{S-S'}) * (Z^\varepsilon_{S'} dM^\varepsilon_{S'})\Big)_\varepsilon + \varepsilon^u\mathscr{C}^{(2)}_T,\end{aligned}$$

which after changing the order of integrations yields

$$N^\varepsilon_T(\phi) = \int_0^T (\phi, Z^\varepsilon_S dM^\varepsilon_S)_\varepsilon + \varepsilon^u\mathscr{C}^{(2)}_T. \tag{4.6}$$

Since $Z^\varepsilon \Rightarrow \mathcal{Z}$, from (4.5) we clearly have that $N^\varepsilon_T(\phi)$ converges weakly to $N_T(\phi)$, which together with (4.6) implies that $N_T(\phi)$ is a local martingale under the law of $\mathcal{Z}$.

Turing to $\Lambda_T(\phi)$ of (4.4), we similarly set

$$\Lambda^\varepsilon_T(\phi) := N^\varepsilon_T(\phi)^2 - \alpha\int_0^T \big(\phi^2, (Z^\varepsilon_S)^2\big)_\varepsilon dS, \tag{4.7}$$

which clearly converges weakly to $\Lambda_T(\phi)$. The quadratic variation of the first term on the RHS of (4.6) is $B^\varepsilon_T(\phi)$. Hence combining (4.6) and (4.7), we have that

$$\Lambda^\varepsilon_T(\phi) = \text{martingle} + \varepsilon^u\mathscr{C}^{(1)}_T + \left(B^\varepsilon_T(\phi) - \alpha\int_0^T \big(\phi^2, (Z^\varepsilon_S)^2\big)_\varepsilon dS\right).$$

Hence, from Proposition 4.1 follows that $\Lambda_T(\phi)$ is a local martingale. □

5. Quadratic Variation

In this section we prove Proposition 4.1. To begin with, we introduce some signed kernel functions

$$(\mathbf{q}_{jkl})_t(x) := \big(\nabla^-\mathbf{p}_t\big)(x+j-k)\big(\nabla^-\mathbf{p}_t\big)(x+j-l), \tag{5.1}$$

$$\mathbf{q}_{ll'} := \sum_{k>|l|}\sum_{j=1\vee(1+l)}^{k\wedge(k+l)} r_k\mathbf{q}_{jkl'}, \tag{5.2}$$

and their total masses

$$\rho_{|k-l|} := \int_0^\infty \sum_x (\mathbf{q}_{jkl})_t(x)dt, \tag{5.3}$$

$$\rho_{ll'} := \int_0^\infty \sum_x (\mathbf{q}_{ll'})_t(x)dt = \sum_{k>|l|} r_k(k-|l|)\rho_{|k-l'|}. \tag{5.4}$$

The integrals exist because of (A.13). Note that although $\mathbf{q}_{jkl}$ depends on (j,k,l), $\rho_{|k-l|}$ depends only on $|k-l|$, because we have summed over x in (5.1).

The proof of Proposition 4.1 relies on a key identity and a key inequality, sated as in Lemma 5.1, which are properties *solely* about the discrete heat kernel $\mathbf{p}$.

Lemma 5.1. *For any* $k \neq 0$,

$$\sum_{k'=1}^{m} \sum_{|l'|<k'} r_{k'}(k'-|l'|)\rho_{|k-l'|} = 0, \tag{5.5}$$

$$\sum_{k'=1}^{m} \Big| \sum_{|l'|<k'} r_{k'}(k'-|l'|)\rho_{|k-l'|} \Big| < 1. \tag{5.6}$$

Lemma 5.1 is proven in Appendix. The identity (5.5) will be used in the proof of Proposition 5.4 and (5.6) will be used in the proof of Lemma 5.7.

Let $\mathscr{F}^\varepsilon_T$ be the canonical filtration of Z^ε_T, that is, $\mathscr{F}^\varepsilon_T := \sigma(Z^\varepsilon_S : S \in [0,T])$. Set

$$(f_l)_t(x) := \varepsilon^{-1} \sum_{k>|l|} r_k \sum_{j=1\vee(1+l)}^{k\wedge(k+l)} \Big(\nabla^- Z^\varepsilon_T\Big)\big(X+j\varepsilon\big)\Big(\nabla^- Z^\varepsilon_T\Big)\big(X+(j-k)\varepsilon\big), \tag{5.7}$$

$$(g_l)_{t,s}(x) := \mathbb{E}\big((f_l)_t(x)\big|\mathscr{F}^\varepsilon_S\big). \tag{5.8}$$

5.1. An integral equation.

In this section we first reduce Proposition 4.1 to the convergence of g_l to zero, that is, Lemma 5.3, and then derive the integral equation (5.16) of g_l. We first need some analysis of the predictable quadratic variation of M^ε.

Proposition 5.2. *$d\langle M^\varepsilon(X), M^\varepsilon(Y)\rangle$ vanishes unless $|X-Y| < m\varepsilon$. Moreover, for $|l| < m$*

$$Z^\varepsilon_T(X)Z^\varepsilon_T(X+l\varepsilon)d\langle M^\varepsilon(X), M^\varepsilon(X+l\varepsilon)\rangle = \Big(\alpha_l Z^\varepsilon_T(X)^2 - (f_l)_t(x) + \varepsilon^{\frac{1}{2}}\mathscr{C}^{(1)}\Big)\frac{dT}{\varepsilon},$$

where

$$\alpha_l := \sum_{k>|l|} r_k(k-|l|). \tag{5.9}$$

Proof. Since the Poisson processes defined in (2.1) are independent, only jumps across *both* the site $x-\frac{1}{2}$ and $x+l-\frac{1}{2}$ contribute to $d\langle M^\varepsilon(X), M^\varepsilon(X+l\varepsilon)\rangle$. When such jumps occur, in view of Figure 5.1, we must have $|l| < m$ and $1 \leq j-l \leq k$. Using (2.5) and the identity

$$d\big\langle P^k(Y), P^k(Y)\big\rangle = d\big\langle Q^k(Y), Q^k(Y)\big\rangle = \varepsilon^{-2}dT,$$

we compute

$$\begin{aligned} &d\langle M^\varepsilon(X), M^\varepsilon(X+l\varepsilon)\rangle \\ &= \mathbb{1}_{\{|l|<m\}} \sum_{k=1}^{m} \sum_{j=1\vee(1+l)}^{k\wedge(k+l)} \Big((e^{-2\lambda}-1)^2 u^{jk}_t(x) + (e^{2\lambda}-1)^2 v^{jk}_t(x)\Big)\varepsilon^{-2}dT. \end{aligned}$$

Further, since $p = \frac{1}{2}(1-\varepsilon^{1/2})$ and $q = \frac{1}{2}(1+\varepsilon^{1/2})$, we have

$$q u^{jk}_t(x) + p v^{jk}_t(x) = \frac{1}{4}\Big(1 - \eta_t(x+j-k)\eta_t(x+j)\Big) + \frac{\varepsilon^{\frac{1}{2}}}{4}\Big(\eta_t(x+j-k) - \eta_t(x+j)\Big).$$

Now, using $e^{2\lambda} = q/p$, the definitions (2.3) and (2.2) of u^{jk} and v^{jk}, and the preceding identity, we obtain

$$Z^\varepsilon_T(X)Z^\varepsilon_T(X+l\varepsilon)d\langle M^\varepsilon(X), M^\varepsilon(X+l\varepsilon)\rangle = \mathbb{1}_{\{|l|<m\}}\Big(D^l_T(X) + \varepsilon^{\frac{1}{2}}E^l_T(X)\Big)\varepsilon^{-1}dT, \tag{5.10}$$



FIGURE 2. Jumps across the the sites $x-\frac{1}{2}$ and $x+l-\frac{1}{2}$.

where

$$D_T^l(X) = Z_T^\varepsilon(X) Z_T^\varepsilon(X+l\varepsilon) \sum_{k>|l|} \sum_{j=1\vee(1+l)}^{k\wedge(k+l)} r_k \Big(1-\eta_t(x+j-k)\eta_t(x+j)\Big)\frac{dT}{4pq}, \tag{5.11}$$

$$E_T^l(X) = Z_T^\varepsilon(X) Z_T^\varepsilon(X+l\varepsilon) \sum_{k>|l|} \sum_{j=1\vee(1+l)}^{k\wedge(k+l)} r_k \Big(\eta_t(x+j-k)-\eta_t(x+j)\Big)\frac{dT}{4pq}.$$

The bound (3.29) for $k=1$ implies that $E_T^l(X)$ is of the form $\mathscr{C}_T^{(1)}(X)$. Similarly, using (2.11) and (3.29) for $k=1$ we know that for all $|j'|,|j''|\leq m$,

$$Z_T^\varepsilon(Y)(Z_T^\varepsilon(Y+j'\varepsilon)-Z_T^\varepsilon(Y+j''\varepsilon)) = \varepsilon^{\frac{1}{2}}\mathscr{C}_T^{(1)}(Y),$$

which implies that in (5.11) we can replace $Z_T^\varepsilon(X)$, $Z_T^\varepsilon(X+l\varepsilon)$ by $Z_T^\varepsilon(X+j'\varepsilon)$, $Z_T^\varepsilon(X+j''\varepsilon)$ for any $|j'|,|j''|\leq m$. Next, using the estimate

$$\eta_t(y)Z_T^\varepsilon(Y) = -\varepsilon^{-\frac{1}{2}}\nabla^- Z_T(Y) + O(\varepsilon^{\frac{1}{2}})Z_T^\varepsilon(Y),$$

(which follows from Talyor expanding (1.5) and the fact $|\lambda|\leq C\varepsilon^{1/2}$) and the identity $4pq=(1-\varepsilon)^{1/2}$ (which follows from (1.6a)), we conclude this proposition. □

Note that $\sum_{|l|<m}\alpha_l$ is equal to α of (1.8). Hence, using Proposition 4.1 and the smoothness of ϕ, we turn $B_T^l(\phi)$, as defined in (4.1), into

$$B_T^l(\phi) = \alpha\int_0^T \big(\phi^2,(Z_S^\varepsilon)^2\big)_\varepsilon dS - F_T^\varepsilon + \varepsilon^{\frac{1}{2}}\mathscr{C}_T^{(1)}, \tag{5.12}$$

where

$$F_T^\varepsilon := \int_0^T \varepsilon\sum_X \phi(X)^2 \sum_{|l|<m} (f_l)_s(x)dS. \tag{5.13}$$

From (5.12), Proposition 4.1 follows as soon as we show

$$\mathbb{E}[(F_T^\varepsilon)^2]\to 0, \text{ uniformly on } T\in[0,\tilde{T}]. \tag{5.14}$$

Next, observe that $\mathbb{E}[(F^{\varepsilon}_T)^2] = 2\mathbb{E}(\int_0^T G^{\varepsilon}_S H^{\varepsilon}_{T,S} dS)$, where

$$G^{\varepsilon}_S := \varepsilon \sum_X \phi(X)^2 \sum_{|l|<m} \big(f_l\big)_s(x),$$

$$H^{\varepsilon}_{T,S} := \varepsilon \sum_X \phi(X)^2 \int_S^T \sum_{|l|<m} \big(g_l\big)_{t',s}(x) dT'. \tag{5.15}$$

Applying the identity (2.11) to the definition (5.7) of f_l, and using the bound (3.29) for $k = 3$, we get $\mathbb{E}(|G^{\varepsilon}_S|^3), \mathbb{E}(|H^{\varepsilon}_{T,S}|^3) \leq C$. In particular, $\mathbb{E}(|H^{\varepsilon}_{T,S}|^2) \leq C$ and $\Big\{(G^{\varepsilon}_T)^2 : \varepsilon > 0, T \in [0, \tilde{T}]\Big\}$ is uniformly integrable. These two properties reduce (5.14) to

Lemma 5.3. $\mathbb{E}\big|H^{\varepsilon}_{T,S}\big| \to 0$ *uniformly on* $\varepsilon^{1/2} \leq S < T \leq \tilde{T}$.

The lower bound $\varepsilon^{1/2}$ is a technical assumption that our latter analysis requires.

We now derive an integral equation of g_l.

Proposition 5.4. *for any* $u \in (0, \frac{1}{2})$*, any* $\varepsilon^{1/2} \leq S < T \leq \tilde{T}$*,*

$$(g_l)_{t,s} = -\int_s^t \sum_{|l'|<m} (\mathbf{q}_{ll'})_{t-t'} \star (g_{l'})_{t',s} dt' + \varepsilon^u (T-S)^{-\frac{1}{2}} \mathscr{C}^{(1)}. \tag{5.16}$$

Proof. First we take discrete gradient of (3.4) to get

$$\begin{aligned}\varepsilon^{-\frac{1}{2}} \nabla^- Z^{\varepsilon}_T = \varepsilon^{-\frac{1}{2}} \nabla^- \mathbf{P}^{\varepsilon}_T * Z^{\varepsilon}_0 + \int_0^T \varepsilon^{-\frac{1}{2}} \nabla^- \mathbf{P}^{\varepsilon}_{T-S} * Z^{\varepsilon}_S dM^{\varepsilon}_S \\ + \varepsilon^{-\frac{1}{2}} \nabla^- \widehat{\mathbf{P}}^{\varepsilon}_{T,0} * Z^{\varepsilon}_0 + \int_0^T \varepsilon^{-\frac{1}{2}} \nabla^- \widehat{\mathbf{P}}^{\varepsilon}_{T,S} * Z^{\varepsilon}_S dM^{\varepsilon}_S.\end{aligned}$$

Let V_T, V'_T, I_T, and I'_T denote the terms on the RHS, respectively. Using discrete integration by parts we have $V_T = \varepsilon^{-1/2} \mathbf{P}^{\varepsilon}_T * \nabla^- Z^{\varepsilon}_0$. From this expression, using (3.21), (2.11), and (1.10a), we conclude $V_T = \mathscr{C}^{(2)}$. Next, computing the L^2-norm of V'_T using Lemma 3.3, (A.12), (3.29), and the readily verified identity

$$\varepsilon \int_0^T (T' \vee \varepsilon^{-2})^{-\frac{3}{2}} dT' \leq C,$$

we conclude $V'_T = \mathscr{C}^{(2)}$. For I and I', we use (3.9), (3.10), (1.10a), (3.29), and Lemma 3.3 to get $I = \varepsilon^{u'} T^{-1/2} \mathscr{C}^{(2)}$ and $I' = \varepsilon^{u'} \mathscr{C}^{(2)}$, where $u' := \frac{1}{2}(u + \frac{1}{2})$. From the preceding estimates, we have

$$\begin{aligned}&\varepsilon^{-1} \nabla^- Z^{\varepsilon}_T(X + j\varepsilon) \nabla^- Z^{\varepsilon}_T(X + (j-k)\varepsilon) \\ &\quad = \Big[\Big(V + V'\Big)_T (X + j\varepsilon)\Big] \Big[\Big(V + V'\Big)_T (X + (j-k)\varepsilon)\Big] + \Big(\varepsilon^{u'} T^{-\frac{1}{2}} + \varepsilon^{2u'} T^{-1}\Big) \mathscr{C}^{(1)},\end{aligned}$$

Further, since $\varepsilon^{1/2} \leq S < T$, we have $(\varepsilon^{u'} T^{-\frac{1}{2}} + \varepsilon^{2u'} T^{-1}) \leq 2\varepsilon^u (T-S)^{-1/2}$. Next, calculating conditional expectation using Proposition 5.2, and summing over j, k, we get

$$\big(g_l\big)_{t,s}(x) = U_{T,S} + L_{T,S} - \int_s^t \sum_{|l'|<m} (\mathbf{q}_{ll'})_{t-t'} \star (g_{l'})_{t',s} dt' + \varepsilon^u (T-S)^{-\frac{1}{2}} \mathscr{C}^{(1)}, \tag{5.17}$$

where, since V' is a martingale in T and V_T is $\mathscr{F}^{\varepsilon}_0$-measurable,

$$L_{T,S} := \int_s^t \sum_{|l'|<m} \alpha_{l'} \ (\mathbf{q}_{ll'})_{t-t'} \star \mathbb{E}(Z^{\varepsilon}_{T'} | \mathscr{F}^{\varepsilon}_S) dt', \tag{5.18}$$

$$U_{T,S}(X) := r_k \sum_{k>|l|} \sum_{j=1\vee(1+l)}^{k\wedge(k+l)} [V_T(X + j\varepsilon) + V'_S(X + j\varepsilon)],$$

$$\times [V_T(X+(j-k)\varepsilon) + V'_S(X+(j-k)\varepsilon)].$$

It remains only to show $L, U = \varepsilon^{1/2}\mathscr{C}^{(1)}(T-S)^{-1/2}$. For U, by Cauchy-Schwartz inequality, we need only to show $V_T, V'_S = \varepsilon^{1/2}(T-S)^{-1/2}\mathscr{C}^{(2)}$. We have already showed in the preceding paragraph that $V_T = \varepsilon^{1/2}\mathscr{C}^{(2)}$. For V'_S, we similarly use (3.21), (2.11), and (1.10a), and

$$\varepsilon \int_S^T (T' \vee \varepsilon^{-2})^{-\frac{3}{2}} dT' \leq \varepsilon (T-S)^{-\frac{1}{2}} C,$$

to get $V'_S = \varepsilon(T-S)^{-1/2}\mathscr{C}^{(2)}$.

Next we turn to L. In the defining equation (5.18) of L, we first divide $\mathbb{E}(Z^{\varepsilon}_{T'}(Y)|\mathscr{F}^{\varepsilon}_S)$ into a sum of $\mathbb{E}(Z^{\varepsilon}_{T'}(Y) - Z^{\varepsilon}_{T'}(X)|\mathscr{F}^{\varepsilon}_S)$, $\mathbb{E}(Z^{\varepsilon}_{T'}(X) - Z^{\varepsilon}_{T}(X)|\mathscr{F}^{\varepsilon}_S)$, and $\mathbb{E}(Z^{\varepsilon}_{T}(X)|\mathscr{F}^{\varepsilon}_S)$. Then we apply the Hölder continuities (3.30) and (3.31) to the first two terms to get

$$L_{T,S}(X) = \mathscr{C}^{(1)} \int_s^t \sum_{|l'|<m} \sum_y (\mathbf{q}_{ll'})_{t-t'}(x-y)\big(|\varepsilon(x-y)|^u + |\varepsilon^2(t-t')|^{\frac{u}{2}} + \varepsilon^u\big) e^{2\varepsilon A_0|x-y|} dt'$$

$$+ \mathbb{E}(Z^{\varepsilon}_T(X)|\mathscr{F}^{\varepsilon}_S) \sum_{|l'|<m} \alpha_{l'} \int_0^{t-s} \sum_y (\mathbf{q}_{ll'})_{t'}(y) dt',$$

where α_l is defined as in (5.9). By (A.14), the first term on the RHS is of the form $\varepsilon^u\mathscr{C}^{(1)}$. We next split the second integral on the RHS into the difference of an integral over $(0,\infty)$ and an integral over $(t-s,\infty)$. By (A.13), the integral of $\sum_y (\mathbf{q}_{ll'})_{t'}(y)$ over $(t-s,\infty)$ is bounded by a constant multiple of $(t-s)^{-\frac{1}{2}} = \varepsilon(T-S)^{-\frac{1}{2}}$. Therefore, by (3.29) we have

$$L_{T,S}(x) = \varepsilon^u (T-S)^{-\frac{1}{2}} \mathscr{C}^{(1)} + \mathbb{E}(Z^{\varepsilon}_T(X)|\mathscr{F}^{\varepsilon}_S) \Big(\sum_{|l'|<m} \rho_{ll'}\alpha_{l'} \Big).$$

where $\rho_{ll'}$ is the total mass of $\mathbf{q}_{ll'}$ as defined in (5.4). From (5.4) (5.9) we have

$$\sum_{|l'|<m} \rho_{ll'}\alpha_{l'} = \sum_{k>|l|} r_k(k-|l|) \left(\sum_{|l'|<m} \sum_{k'>|l'|} r_{k'}(k'-|l'|)\rho_{|k-l'|} \right).$$

After interchanging the order of summation in the parentheses, by (5.5) we know that the quantity in the parentheses is zero. □

Remark 5.5. In the case of simple exclusion $m=1$, [3] showed that

$$\int_0^{\infty} \sum_x |(\mathbf{q}_{00})_t(x)| dt < 1, \tag{5.19}$$

which implies the integral operator in (5.16) is a contraction map. (Note that when $m=1$ the index l and l' take only the value 0.) In view of Proposition A.3, (5.19) does not generalize to the case of $m>1$. We circumvent this problem by first reducing (5.16) to a finite dimensional linear equation (Proposition 5.6), and showing contractivity for the finite dimensional equation 5.7.

5.2. **Reduction to finite dimensional equation.**

We first reduce (5.16) to a finite dimensional linear equation.

Proposition 5.6. *Define*

$$(\gamma_l)_{t,s}(x) := \int_S^T \varepsilon^{\frac{3}{4}} \sum_{|y-x|<\varepsilon^{-3/4}} (g_l)_{t',s}(y)\ dT'. \tag{5.20}$$

For all $\varepsilon^{1/2} \leq S < T \leq \tilde{T}$, $u \in (0, \frac{1}{2})$,

$$(\gamma_l)_{t,s}(x) = -\sum_{|l'|<m} \rho_{ll'}\ \big((\gamma_{l'})_{t,s}(x)\big) + \varepsilon^u\mathscr{C}^{(1)}. \tag{5.21}$$



(a) Ranges of summation.

(b) Sub-domains of time integral.

FIGURE 3. Sub-domains.

Proof. First we sum over $y \in (x-\varepsilon^{-3/4}, x+\varepsilon^{-3/4})$, and integrate over (S,T) in (5.16). Subsequently, the LHS becomes $(\gamma_l)_{t,s}(x)$, and the second term on the RHS becomes $\varepsilon^u\mathscr{C}^{(1)}$. Therefore, it remains only to show that the first term on the RHS

$$\sum_{|l'|<m}\left[\int_S^T \varepsilon^{\frac{3}{4}} \sum_{|y-x|<\varepsilon^{-3/4}} \int_s^{t'} \Big((\mathbf{q}_{ll'})_{t-t'} \star (g_{l'})_{t',s}\Big)(y)dt''dT'\right] \tag{5.22}$$

is equal to the sum of $\sum_{|l'|}\rho_{ll'}\ (\gamma_{l'})_{t,s}(x)$ and some error terms of the form $\varepsilon^{1/2}\mathscr{C}^{(1)}$.

After a change of variable in space and a change of variable in time, the term sitting in the bracket in (5.22) becomes

$$\varepsilon^2 \int_{\tilde{\Delta}_s^t} \varepsilon^{\frac{3}{4}} \sum_{(y_1,y_2)\in\tilde{\Delta}_x} \Big[\big(\mathbf{q}_{ll'}\big)_{t_1}(y_1)\Big]\Big[\big(g_{l'}\big)_{t_2+s,s}(y_2)\Big]\, dt_1dt_2, \tag{5.23}$$

where

$$\begin{aligned}\tilde{\Delta}_s^t &= \big\{(t_1,t_2)\in(0,\infty)^2 : t_1+t_2<t-s\big\},\\ \tilde{\Delta}_x &= \Big\{(y_1,y_2)\in\mathbb{Z}^2 : |y_1+y_2-x|<\varepsilon^{-3/4}\Big\}.\end{aligned}$$

We divide the sum into four pieces: $\varepsilon^{\frac{3}{4}}\sum_{\tilde{\Delta}_x} = \varepsilon^{\frac{3}{4}}\left(\sum_{\mathfrak{A}}+\sum_{\mathfrak{B}}+\sum_{\mathfrak{C}}-\sum_{\mathfrak{C}'}\right)$, where

$$\begin{aligned}\mathfrak{A} &:= \Big\{|y_1|<\varepsilon^{-\frac{1}{4}}\Big\}\times\Big\{|y_2-x|<\varepsilon^{-\frac{3}{4}}\Big\}, & \mathfrak{B} &:= \tilde{\Delta}_x\cap\Big\{|y_1|\geq\varepsilon^{-\frac{1}{4}}\Big\},\\ \mathfrak{C} &:= \Big(\tilde{\Delta}_x\cap\Big\{|y_1|<\varepsilon^{-\frac{1}{4}}\Big\}\Big)\setminus\mathfrak{A}, & \mathfrak{C}' &:= \mathfrak{A}\setminus\Big(\tilde{\Delta}_x\cap\Big\{|y_1|<\varepsilon^{-\frac{1}{4}}\Big\}\Big).\end{aligned}$$

See Figure 3(a). The sums $\sum_{\mathfrak{B}}$, $\sum_{\mathfrak{C}}$, and $\sum_{\mathfrak{C}'}$ have negligible contributions, as we next show. By the definition (5.7) of f_l, and the estimates (2.11) and (3.29),

$$\big\|(g_l)_{t_2,s}(x)\big\|_1 \leq Ce^{2A_0\varepsilon|x|}. \tag{5.24}$$

By (5.24) and (A.13), the sums over $\mathfrak{C}$ and $\mathfrak{C}'$ are of the form $\varepsilon^{3/4-1/4}(1\wedge t_1^{-3/2})\mathscr{C}^{(1)}$. Next, by (5.24) the sum over $\mathfrak{B}$ is of the form $\sum\limits_{|y_1|\geq\varepsilon^{-1/4}}|(\mathbf{q}_{ll'})_{t_1}(y_1)|\mathscr{C}^{(1)}$. Using (A.13) for $\varepsilon=A=1$ we have

$$\sum_{y_1}\Big|\big(\mathbf{q}_{ll'}\big)_{t_1}(y_1)\Big|e^{|x|}\leq\Big(t_1^{-\frac{3}{2}}\wedge 1\Big). \tag{5.25}$$

Therefore, the sum over $\mathfrak{B}$ is of the form $e^{-\varepsilon^{-1/4}}(t_1^{-3/2}\wedge 1)\mathscr{C}^{(1)}$. The sum over $\mathfrak{A}$ reads

$$\Bigg[\sum_{|y_1|<\varepsilon^{-1/4}}(\mathbf{q}_{ll'})_{t-t'}(y)\Bigg]\Bigg[\varepsilon^{\frac{3}{4}}\sum_{|y-x|<\varepsilon^{-3/4}}(g_{l'})_{t_2+s,s}(y)\Bigg].$$

Using (5.24) and (5.25), we turn the preceding sum over $|y_1|<\varepsilon^{-1/4}$ into a sum over $\mathbb{Z}$ and a negligible error:

$$\Bigg[\sum_{y\in\mathbb{Z}}(\mathbf{q}_{ll'})_{t-t'}(y)\Bigg]\Bigg[\varepsilon^{\frac{3}{4}}\sum_{|y-x|<\varepsilon^{-3/4}}(g_{l'})_{t_2+s,s}(y)\Bigg]+\Big(1\wedge t_1^{-\frac{3}{2}}\Big)e^{-A\varepsilon^{-\frac{1}{4}}}\mathscr{C}^{(1)}.$$

Therefore,

$$\begin{aligned}&\varepsilon^{\frac{3}{4}}\sum_{(y_1,y_2)\in\tilde{\Delta}_x}\Big[\big(\mathbf{q}_{ll'}\big)_{t_1}(y_1)\Big]\Big[\big(g_{l'}\big)_{t_2+s,s}(y_2)\Big]\\&=\Bigg[\sum_{y}(\mathbf{q}_{ll'})_{t_1}(y)\Bigg]\Bigg[\varepsilon^{\frac{3}{4}}\sum_{|y-x|<\varepsilon^{-3/4}}(g_{l'})_{t_2+s,s}(y)\Bigg]+w,\end{aligned}$$

where

$$w=\Big(\varepsilon^{\frac{1}{2}}+e^{-A\varepsilon^{-1/4}}\Big)\Big(1\wedge t_1^{-\frac{3}{2}}\Big)\mathscr{C}^{(1)}. \tag{5.26}$$

Next we integrate over $\tilde{\Delta}^t_s$. The time integral of the coefficient multiplying $\mathscr{C}^{(1)}$ in (5.26)

$$\varepsilon^2\int_{\tilde{\Delta}^t_s}\Big(\varepsilon^{\frac{1}{2}}+e^{-A\varepsilon^{-1/4}}\Big)\Big(1\wedge t_1^{-\frac{3}{2}}\Big)dt_1dt_2$$

is less than or equal to a constant multiple of $\varepsilon^{1/2}$. Hence (5.23) is equal to

$$\varepsilon^2\int\limits_{\tilde{\Delta}^t_s}\Bigg[\sum_{y}(\mathbf{q}_{ll'})_{t_1}(y)\Bigg]\Bigg[\varepsilon^{\frac{3}{4}}\sum_{|y-x|<\varepsilon^{-3/4}}(g_{l'})_{t_2+s,s}(y)\Bigg]dt_1dt_2+\varepsilon^{\frac{1}{2}}\mathscr{C}^{(1)}.$$

For the integral over $\tilde{\Delta}^t_s$, we use the same type of argument as before. First we divide the integral over $\tilde{\Delta}^t_s$ as $\varepsilon^2\int_{\tilde{\Delta}^t_s}=\varepsilon^2\big(\int_{\mathfrak{A}}+\int_{\mathfrak{B}}-\int_{\mathfrak{C}}\big)$, where

$$\tilde{\mathfrak{A}}:=\big\{0<t_1<\varepsilon^{-1}\big\}\times\{0<t_2<t-s\},\quad\tilde{\mathfrak{B}}:=\tilde{\Delta}^t_s\cap\big\{t_1\geq\varepsilon^{-1}\big\},\quad\tilde{\mathfrak{C}}:=\tilde{\mathfrak{A}}\setminus\tilde{\Delta}^t_s.$$

See Figure 3(b). Using (A.13) for $A=0$ and (5.24) we get that the integral over $\tilde{\mathfrak{A}}$ is $\rho_{ll'}(\gamma_{l'})_{t,s}(x)+\varepsilon^{\frac{1}{2}}\mathscr{C}^{(1)}$. Similarly, by (A.13) for $A=0$ and (5.25) the integral over $\tilde{\mathfrak{B}}$ is of the form $\varepsilon^{\frac{1}{2}}\mathscr{C}^{(1)}$, and the integral over $\tilde{\mathfrak{C}}$ is of the form $\varepsilon^{2-1}\mathscr{C}^{(1)}$. □

Next we show that the matrix ρ appearing in (5.21) has spectrum radius less than one.

Lemma 5.7. *Let ρ denote the $(2m-1)\times(2m-1)$ matrix with entry $\rho_{ll'}$. All eigenvalues of ρ have absolute values strictly less than one. In particular, $I+\rho$ is invertible.*

Proof. By Gelfand's formula $\sigma(\rho) = \lim_{n\to\infty} \|\rho^n\|^{1/n}$, which holds for any norm $\|\cdot\|$ on finite dimensional space, we need only to show $\lim_{n\to\infty} \rho^n = 0$ (since then we will have $\|\rho^{n_0}\| \leq 1/2$ for some n_0). let a denote the $(2m-1)\times m$ matrix with entry $a_{lk} = \rho_{|k-l|}$ and b denote the $m \times (2m-1)$ matrix with entry $b_{kl} = r_k(k-|l|)_+$, where the subscript $+$ refers to the the positive part. From (5.4) we know $\rho = ab$. The statement $\lim_{n\to\infty}(ab)^n = 0$ is then equivalent to $\lim_{n\to\infty}(ba)^n = 0$. Indeed,

$$\|ba\|_{L^\infty\to L^\infty} = \sum_{k'=1}^{m} \Big| \sum_{|l'|<k'} r_{k'}(k'-|l'|)\rho_{|k-l'|} \Big|,$$

which by (5.6) is strictly less than one. □

Remark 5.8. From Lemma 5.7 and Proposition 5.6 the bound

$$(\gamma_l)_{t,s} = \varepsilon^u \mathscr{C}^{(1)}, \qquad \text{whenever } \varepsilon^{\frac{1}{2}} \leq S < T \leq \tilde{T} \tag{5.27}$$

follows at once. Recall that γ_l (5.20) is a space-time average of g_l, which at first sight seems useless to prove Lemma 5.3. However, by the smoothness of ϕ, at microscopic level ϕ remains a constant. Therefore, the sum in (5.15) behaves like an average over space of g_l.

We now implement this idea to give a proof of Lemma 5.3.

Proof of Lemma 5.3. Let $[-L, L]$ be a macroscopic interval off which ϕ vanishes. Set

$$\begin{aligned} x_i^\varepsilon &:= \lfloor 2i\varepsilon^{-3/4} \rfloor, \quad I_i^\varepsilon := \Big(-\varepsilon^{-\frac{3}{4}} + x_i^\varepsilon, x_i^\varepsilon + \varepsilon^{-\frac{3}{4}}\Big) \cap \mathbb{Z}, \\ \mathfrak{X}^\varepsilon &:= \big\{ i \in \mathbb{Z} : x_i^\varepsilon \in [-\varepsilon^{-1}L, \varepsilon^{-1}L] \big\}, \\ \mathfrak{E}^\varepsilon &:= \Big\{ x_i^\varepsilon \pm \varepsilon^{-\frac{3}{4}} : i \in \mathfrak{X}^\varepsilon \Big\} \cap \mathbb{Z}, \end{aligned}$$

Note that $\#\mathfrak{X}^\varepsilon, \#\mathfrak{E}^\varepsilon \leq \varepsilon^{-1/4}C$. We then divide the sum (5.15) into sub-sums as

$$\begin{aligned} &\varepsilon^{\frac{1}{4}} \sum_{i\in\mathfrak{X}^\varepsilon} \left(\varepsilon^{\frac{3}{4}} \sum_{y\in I_i^\varepsilon} \phi(Y)^2 \int_{\varepsilon^{-2}S}^{\varepsilon^{-2}T} \sum_{|l|<m} (g_l)_{t',s}(y) dT' \right) \\ &\quad + \varepsilon \sum_{y\in\mathfrak{E}^\varepsilon} \phi(Y)^2 \int_{\varepsilon^{-2}S}^{\varepsilon^{-2}T} \sum_{|l|<m} (g_l)_{t',s}(y) dT'. \end{aligned} \tag{5.28}$$

Since $\#\mathfrak{E}^\varepsilon \leq \varepsilon^{-1/4}C$, by (5.24) the second term of (5.28) is of the form $\varepsilon^{-3/4}\mathscr{C}^{(1)}$, which is negligible. On each subinterval I_i^ε we replace $\phi(Y)$ by $\phi(\varepsilon^{-1}x_i^\varepsilon)$, at the cause of an error term whose L^1-norm is bounded by $\|\phi'\|_\infty \varepsilon^{1/4}$. This replacement turns the first term of (5.28) into

$$\varepsilon^{\frac{1}{4}} \sum_{i\in\mathfrak{X}^\varepsilon} \phi(\varepsilon^{-1}x_i^\varepsilon)^2 (\gamma_i)_{t,s}(x) + \varepsilon^{\frac{1}{4}}\mathscr{C}^{(1)}.$$

Lemma 5.3 now follows directly from (5.27). □

6. EXACT STATISTICS.

In this section we prove Theorem 1.3 and 1.4. In view of (1.4) and (1.6), for the step initial condition (1.12) we set

$$Z_0^\varepsilon(X) = \frac{\varepsilon^{-\frac{1}{2}}}{2} \exp\left(-\lambda_\varepsilon |X|\right) \tag{6.1}$$

and for the step Bernoulli initial condition (1.13) we set

$$Z_0^\varepsilon(X) = \left\{ \begin{array}{ll} \exp\left(-\varepsilon^{-1}\lambda_\varepsilon|X|\right) & \text{, when } X \leq 0, \\ \exp\left(-\varepsilon^{-1}\lambda_\varepsilon \mathrm{RW}_{|\varepsilon^{-1}X|}\right) & \text{, when } X > 0, \end{array} \right. \tag{6.2}$$

where λ_ε is defined as in (1.6b), and RW_n stands for the n-steps position of a symmetric simple random walk on $\mathbb{Z}$, starting at 0. Note the extra factor of $\varepsilon^{-\frac{1}{2}}/2$ in (6.1), which ensures that Z^ε_0 converges to a non-zero limit. Theorem 1.3 and 1.4 is proven by apply Theorem 1.1, and the following two Propositions.

Proposition 6.1. *Consider the initial conditions* (6.1) *and* (6.2). *Given any* $k \in \mathbb{N}$, $\delta > 0$, *there exist* $A \geq 0$, $C(\delta) \geq 0$ *such that*

$$\|Z^\varepsilon_\delta(X)\|_k \leq C(\delta) e^{A|X|}, \tag{6.3}$$

$$\|Z^\varepsilon_\delta(X) - Z^\varepsilon_\delta(X')\|_k \leq |X - X'|^u e^{A(|X|+|X'|)} C(\delta). \tag{6.4}$$

Proposition 6.2. *Given any* C^∞*-function* ϕ *with compact support, for the initial conditions* (6.1) *and* (6.2), *any* $T \in (0, \tilde{T}]$,

$$\int_{\mathbb{R}} \|\mathcal{Z}_T(Y)\|_2 |\phi(Y)| dY \leq C, \tag{6.5}$$

$$\int_{\mathbb{R}} P_{T-\delta}(X - Y)\mathcal{Z}_{\alpha^{-1}\delta}(Y) dY \Rightarrow \int_{\mathbb{R}} P_T(X - Y)\mathcal{Z}_{in}(Y) dY, \text{ as } \delta \to 0. \tag{6.6}$$

Indeed, $Z^\varepsilon_0(\,\cdot\,)$ converges weakly to $\mathcal{Z}_{\mathrm{in}}(\,\cdot\,)$, where

$$\mathcal{Z}_{\mathrm{in}}(X) = \begin{cases} \delta_0(X) & \text{, for (6.1)}, \\ \mathbb{1}_{(0,\infty)}(X) B(X), & \text{for (6.2)}, \end{cases} \tag{6.7}$$

δ_0 the delta measure concentrated at 0, and $B(\,\cdot\,)$ is the standard Brownian motion. Since the one point distribution of solutions to the SHE (1.11) for the initial conditions (6.7) have been derived by [1] and [7], we need only to show convergence of Z^ε to the solution of the SHE. However, Theorem 1.1 does not apply directly, because (6.1) violates (1.10a) and (1.10b), and (6.2) violates (1.10b). We circumvent this problem following the same argument of [1], by first showing that the conditions (1.10a) and (1.10b) hold for Z^ε_δ, for any $\delta > 0$, namely Proposition 6.1, and applying Theorem 1.1 to conclude $Z^\varepsilon_\cdot(\,\cdot\,) \Rightarrow \mathcal{Z}_\cdot(\,\cdot\,)$ on $[\delta, \infty) \times \mathbb{R}$, for some $\mathcal{Z}$ satisfying the SHE (1.11) on $[\delta, \infty) \times \mathbb{R}$. Further, the extension argument in [1, Section 3] (which applies to generic processes satisfying (1.11)) extends $\mathcal{Z}$ to a process defined on $(0, \infty) \times \mathbb{R}$, yielding $Z^\varepsilon_\cdot(\,\cdot\,) \Rightarrow \mathcal{Z}_\cdot(\,\cdot\,)$ on $(0, \infty) \times \mathbb{R}$, and

$$\mathcal{Z}_{\alpha^{-1}T} = \int_{\mathbb{R}} P_T(X - Y)\mathcal{Z}_{\alpha^{-1}\delta}(Y) dY + \int_{[\delta,T]} \int_{\mathbb{R}} P_{T-S}(X - Y)\mathcal{Z}_{\alpha^{-1}S}(Y) W(dY, dS),$$

for any $\delta > 0$. Finally, the estimates (6.6) and (6.5) allow to pass to take limit $\delta \to 0$ of the first and second terms on the RHS, respectively, and this shows that $\mathcal{Z}$ satisfies the SHE (1.11) with initial condition $\mathcal{Z}_{\mathrm{in}}$.

From the preceding discussion, we need only to prove Propositions 6.1 and 6.2. We first establish a technical Lemma.

Lemma 6.3. *There exists a positive constant* C *such that* $|\tilde{\mathbf{P}}^\varepsilon_{T,S}|'_0 \leq C$, *for all* $0 \leq S < T \leq \tilde{T}$, *where the norm* $|\cdot|'_0$ *is defined as*

$$|F|'_0 := \sup_X \varepsilon \sum_Y \|F(X, Y)\|_\infty. \tag{6.8}$$

Proof. Since (3.16)–(3.19) are symmetric in X and Y, the proof of (3.22) also shows

$$e^{-A|Y|}\varepsilon \sum_X \Big\| \tilde{\mathbf{P}}^\varepsilon_{T,S}(X, Y) - \mathbf{P}^\varepsilon_{T-S}(X - Y) \Big\|_\infty e^{A|X|} \leq C\varepsilon^u,$$

which concludes the lemma. ☐

Proof of Proposition 6.1. Consider first the case of (6.2). Indeed, (1.10a) holds for any $k \in \mathbb{N}$, $A > 0$. Hence, by (3.29)

$$\|Z^\varepsilon_T(X)\|_k \leq e^{|X|} C, \tag{6.9}$$

which implies (6.3). Next, observe that the assumption (1.10b) is used in the proof of (1.10b) only in (3.33). We then replace (3.33) by

$$\big\|\nabla_n^- \mathbf{P}^\varepsilon_\delta Z^\varepsilon(X,X')\big\|_k \le \Big(\varepsilon\sum_Y \big\|\nabla_n^-\mathbf{P}^\varepsilon_\delta(X-Y)\big\|_\infty^2 \|Z_0^\varepsilon(Y)\|_k^2 e^{|Y|}\Big)^{\frac12}\Big(\varepsilon\sum_Y e^{-|Y|}\Big)^{\frac12},$$

and then use (6.3), (A.6) for $u = 1$, and (A.10) to bound the RHS by a constant multiple of $C\delta^{-1}|X-X'|^{\frac12}$. This proves (6.4).

Next consider the case of (6.1), using Lemma 3.3, (3.4), and iteration, we get, for any $k \in \mathbb{N}$,

$$\big\|(Z^\varepsilon_\delta)^2\big\|_k \le C\,E_\delta + \sum_{n=1}^{\infty} C^n\,B^n_\delta, \tag{6.10}$$

where

$$E_\delta := \Big(\|\tilde{\mathbf{P}}^\varepsilon_{\delta,0}\|_\infty * (Z^\varepsilon_0)\Big)^2, \quad B^n_\delta := \int_{\Delta^\delta_0(n)} D_{\delta,T_1} * \cdots * D_{T_{n-1},T_n} * E_{T_n}\,dT_1\cdots dT_n,$$
$$D_{T,S} := \mathcal{F}(\tilde{\mathbf{P}}^\varepsilon_{T,\cdot})_S,$$

and $\Delta^\delta_0(n)$ is defined as in (3.12). For any random field $G(Y)$, define the norm

$$|G|''_0 := \varepsilon\sum_Y \|G(Y)\|_\infty. \tag{6.11}$$

By definition,

$$|Z^\varepsilon_0|''_0 \le C, \tag{6.12}$$

which together with (A.4) imply

$$E_T \le CT^{-\frac12}(\|\tilde{\mathbf{P}}^\varepsilon_{T,0}\|_\infty * Z^\varepsilon_0), \tag{6.13}$$
$$E_T \le CT^{-1}. \tag{6.14}$$

By definition given any $(T_0,\ldots,T_n) \in \Delta^T_0(n)$, for some l we must have $T_l - T_{l+1} \ge T/(n+1)$. Consequently applying (A.4) and (6.13) to the definition of B^n, we obtain

$$B^n_T \le C\sum_{l=0}^{n}\int_{\Delta^\delta_0(n)}\Big(\frac{n+1}{T}\Big) I_l\, I'_l\, dT_1\cdots dT_n, \tag{6.15}$$

where

$$I_l := \prod_{i=0}^{l-1}(T_i-T_{i+1})^{-\frac12}\Big|\tilde{\mathbf{P}}^\varepsilon_{T_i,T_{i+1}}\Big|_0, \quad I'_l := \Big(\prod_{i=l+1}^{n}(T_i-T_{i+1})^{-\frac12}\Big|\tilde{\mathbf{P}}^\varepsilon_{T_i,T_{i+1}}\Big|'_0\Big)|Z^\varepsilon_0|''_0.$$

Here the norms $|\cdot|_0$, $|\cdot|'_0$, and $|\cdot|''_0$ are defined as in (3.5), (6.8) , and (6.11), respectively. Applying Lemma 6.3 to (6.15), and evaluation the multiple time integral, we get, for any $\beta \in (0,\frac12)$, $B^n_T \le n^2C^n[T^n/(n!)]^\beta$, which together with (6.14) proves (6.3).

Turning to (6.4), set $j := \varepsilon^{-1}(X'-X) \in \mathbb{Z}$. From (3.4) and Lemma 3.3, we have, for any $k \in \mathbb{N}$,

$$\big\|(\nabla_j^- Z^\varepsilon_\delta)^2\big\|_k \le C\Big(\|\nabla_j^-\tilde{\mathbf{P}}^\varepsilon_{\delta,0}\|_\infty * Z^\varepsilon_0\Big)^2 + C\int_0^T \mathcal{F}\Big(\nabla_j^-\tilde{\mathbf{P}}^\varepsilon_{T,\cdot}\Big)_S * \big\|(Z^\varepsilon_S)^2\big\|_k dS.$$

Using (3.23) and (3.24), for any $u \in (0,\frac12)$ we bound the first term on the RHS by a constant multiple of $\delta^{-(1+u)}|X-X'|^{2u}$. Next, we plug (6.10) into the second term on the RHS, and use (3.24) and the same argument proving (6.15) to bound the second term on the RHS by

$$|X-X'|^{2u}\sum_{n=1}^{\infty} C^n\,B'^n_T,$$

where

$$B'^{n}_{T} := \sum_{l=0}^{n} \int_{\Delta_0^\delta(n)} \Big(\frac{n+1}{T}\Big) J_l\, J'_l\, dT_1 \cdots dT_n,$$

$$J_l := \prod_{i=0}^{l-1} (T_i - T_{i+1})^{-(\frac{1}{2}+u)} \Big|\tilde{\mathbf{P}}^{\varepsilon}_{T_i,T_{i+1}}\Big|_0, \quad J'_l := \Big(\prod_{i=l+1}^{n} (T_i - T_{i+1})^{-(\frac{1}{2}+u)} \Big|\tilde{\mathbf{P}}^{\varepsilon}_{T_i,T_{i+1}}\Big|'_0 \Big) |Z^{\varepsilon}_0|''_0.$$

Finally, applying Lemma 6.3 , and evaluation the multiple time integral, we get, for any $\beta \in (0, \frac{1}{2} - u)$, $B'^{n}_{T} \leq n^2 C^n [T^n/(n!)]^\beta$, which proves (6.4). □

Proof of Proposition 6.2. First we assume (6.5) and prove (6.6). Observe that by (3.4) we have

$$\int_{\mathbb{R}} P_{T-\delta}(X-Y)(I^{\varepsilon}_{\delta} + J^{\varepsilon}_{\delta}) dY \Rightarrow \int_{\mathbb{R}} P_{T-\delta}(X-Y) \mathcal{Z}_{\alpha^{-1}\delta}(Y) dY, \text{ as } \varepsilon \to 0,$$

where

$$I^{\varepsilon}_{\delta} := \tilde{\mathbf{P}}^{\varepsilon}_{\alpha^{-1}\delta,0} * Z^{\varepsilon}_{\text{in}}, \quad J^{\varepsilon}_{\delta}(Y) := \int_0^\delta \tilde{\mathbf{P}}^{\varepsilon}_{\alpha^{-1}\delta,S} * Z^{\varepsilon}_S dM^{\varepsilon}_S.$$

Using (3.9) and (A.1) one can show that

$$\int_{\mathbb{R}} P_{T-\delta}(X-Y) I^{\varepsilon}_{\delta}(Y) dY \Rightarrow \int_{\mathbb{R}} P_T(X-Y) Z_{\text{in}}(Y) dY, \text{ as } \varepsilon \to 0,$$

which implies that J^{ε}_{δ} also convergence to a limit $\mathcal{J}_\delta$ as $\varepsilon \to 0$. By Fatou's lemma,

$$\Big\| \int_{\mathbb{R}} \mathcal{J}_\delta(X) \phi(X) dX \Big\|_2 \leq \liminf_{\varepsilon \to 0} \int_0^\delta \varepsilon^2 \sum_{X,Y} |\phi(X)| \Big\| \tilde{\mathbf{P}}^{\varepsilon}_{\alpha^{-1}\delta,S}(X,Y) \Big\|_\infty \| Z^{\varepsilon}_S(Y) \|_2 dS,$$

which by (6.5) and (3.22) is bounded by a constant multiple of δ. Therefore, $\mathcal{J}_\delta \Rightarrow 0$ as $\delta \to 0$, and this proves (6.6).

Next we turn to (6.5). For the case of (6.2), (6.5) follows directly from (6.9). Consider next the case of (6.1). Using Fatou's lemma, (3.4), Lemma 3.3, (3.23), and the readily verified identity

$$\varepsilon \sum_X \| (F * G)(X) \|_\infty \leq |F|'_0 \, |G|''_0, \tag{6.16}$$

we have

$$\int_{\mathbb{R}} |\phi(X)| \| \mathcal{Z}_T(X) \|_2 dX \leq \liminf_{\varepsilon \to 0} \|\phi\|_\infty \Big\| \varepsilon \sum_X Z^{\varepsilon}_S(Y) \Big\|_2 \leq \liminf_{\varepsilon \to 0} \|\phi\|_\infty (U^\varepsilon + C\, V^\varepsilon),$$

where

$$U^\varepsilon := |\tilde{\mathbf{P}}^{\varepsilon}_{T,0}|'_0 \, |Z^{\varepsilon}_{\text{in}}|''_0, \quad V^\varepsilon := \Big(\int_0^T (T-S)^{-\frac{1}{2}} |\tilde{\mathbf{P}}^{\varepsilon}_{T,S}|'_0 \, |(Z^{\varepsilon}_S)^2|''_0 \, dS \Big)^{\frac{1}{2}}.$$

By (6.12) and Lemma 6.3, $U^\varepsilon \leq C$. Next, by (6.13), (6.10), (3.23), and (6.16), we have

$$\begin{aligned} |(Z^{\varepsilon}_S)^2|''_0 &\leq C S^{-\frac{1}{2}} |\tilde{\mathbf{P}}^{\varepsilon}_{S,0}|'_0 \, |Z^{\varepsilon}_{\text{in}}|''_0 \\ &\quad + \sum_{n=1}^\infty C^n \int_{\Delta_0^S(n)} \Big(\prod_{i=0}^{n} (T_i - T_{i+1})^{-\frac{1}{2}} |\tilde{\mathbf{P}}^{\varepsilon}_{T_i,T_{i+1}}|'_0 \Big) |Z^{\varepsilon}_0|''_0 dT_1 \cdots dT_n, . \end{aligned}$$

Plugging this inequality back to V^ε, using (6.12) and Lemma 6.3, and evaluating the multiple time integral, we show that $V^\varepsilon \leq C$. □

Appendix A. Properties of the discrete heat kernel.

We provide some expressions and estimates of $\mathbf{P}^\varepsilon$, as defined in (3.2). Recall the well-known expressions

$$\mathbf{P}^\varepsilon_T(X) = \frac{1}{2\pi}\int_{-\varepsilon^{-1}\pi}^{\varepsilon^{-1}\pi} e^{-iX\zeta}\exp\big[-\varepsilon^{-2}T(1-\phi(\varepsilon\zeta))\big]d\zeta, \tag{A.1}$$

$$\mathbf{p}_t(x) = e^{-t}\sum_{n=0}^{\infty}\frac{t^n}{n!}\mathbb{P}(\Xi_n = x), \tag{A.2}$$

see [3, Section 5.A], where $\Xi_n = \xi_1 + \ldots + \xi_n$ is the sum of i.i.d. integer-valued random variables with distribution $\mathbb{P}(\xi_1 = k) = \mathbb{P}(\xi_1 = -k) = \frac{r_k}{2}$, and

$$\phi(\zeta) := \mathbb{E}(e^{i\xi_1\zeta}) = \sum_{k=1}^{m} r_k\cos(k\zeta)$$

is the Fourier transform of ξ_1. From the assumption (1.9) follows

$$\inf_{\zeta\in[-\pi,\pi]}\frac{1-\phi(\zeta)}{\zeta^2} > 0. \tag{A.3}$$

Proposition A.1. *Given any $u \in [0,1]$, $A \geq 0$, we have the following L^∞-type estimates of $\mathbf{P}^\varepsilon$:*

$$\sup_X \mathbf{P}^\varepsilon_T(X) \leq CT^{-\frac{1}{2}}, \tag{A.4}$$

$$\sup_X \varepsilon^{-1}\big|\nabla^-\mathbf{P}^\varepsilon_T(X)\big| \leq \big(T^{-1}\wedge\varepsilon^{-2}\big)C, \tag{A.5}$$

$$\sup_{X\neq Y\in\varepsilon\mathbb{Z}}\Big(|X-Y|^{-u}|\mathbf{P}^\varepsilon_T(X)-\mathbf{P}^\varepsilon_T(Y)|\Big) \leq CT^{-\frac{u+1}{2}}, \tag{A.6}$$

$$\sup_X \varepsilon^{-2}|\Delta\mathbf{P}^\varepsilon_T(X)| \leq \Big(T^{-\frac{3}{2}}\wedge\varepsilon^{-3}\Big)C, \tag{A.7}$$

$$\sup_X |\mathbf{P}^\varepsilon_T(X)-\mathbf{P}^\varepsilon_{T'}(X)| \leq CT^{-\frac{1}{2}}|(T-T')/T|^{u/2}, \tag{A.8}$$

$$\varepsilon^{-2}|\Delta\mathbf{P}^\varepsilon_T(X)| \leq C|X|^{-1}\big(T^{-1}\wedge\varepsilon^{-2}\big), \tag{A.9}$$

and the following L^1-type estimates:

$$\varepsilon\sum_X \mathbf{P}^\varepsilon_T(X)e^{A|X|} \leq C, \tag{A.10}$$

$$\varepsilon\sum_X \mathbf{P}^\varepsilon_T(X)|X|^u e^{A|X|} \leq C|T\vee\varepsilon^2|^{\frac{u}{2}}. \tag{A.11}$$

Proof. To bound $\mathbf{P}^\varepsilon$, due to (A.3) we can replace the exponential in (A.1) by $e^{-c\zeta^2}$ for some $c > 0$, and get (A.4) through the bound $|e^{-iX\zeta}| \leq 1$ and integrating over $\zeta \in \mathbb{R}$. Similarly, modifying (A.1) by taking discrete gradient, difference, and discrete Laplacian, we get the bounds (A.5)-(A.8) by applying the same argument, where the extra factor of $\varepsilon^{-\beta}$ follows from (3.2) and the fact that $0 \leq \mathbf{p}_t(x) \leq 1$. The bound (A.9) is derived similarly, by first taking discrete Laplacian to (A.1), integrating by parts with

$$u = e^{-iX\zeta}, \quad dv = 2\big(\cos(\varepsilon\zeta)-1\big)\exp\big[-\varepsilon^{-2}T(1-\phi(\varepsilon\zeta))\big]d\zeta,$$

and applying the bounds (1.9), $\sup_{\mathbb{R}\setminus\{0\}}|\zeta^{-2}(\cos(\zeta)-1)| < \infty$, and $\sup_{\mathbb{R}\setminus\{0\}}|\zeta^{-1}\sin(\zeta)| < \infty$.

From (A.2), we know

$$\varepsilon\sum_X \mathbf{P}^\varepsilon_T(X)e^{A|X|(T\vee\varepsilon^{-2})^{-1/2}} = \sum_x \mathbf{p}_t(x)e^{A|x|(t\vee 1)^{-1/2}} = e^{-t}\sum_{n=0}^{\infty}\frac{t^n}{n!}\mathbb{E}\Big(e^{A|\Xi_n|(t\vee 1)^{-1/2}}\Big).$$

Using triangle inequality $|\Xi_n| \leq |\xi_1| + \ldots + |\xi_n|$ and the fact that $\xi_1, \ldots, \xi_n$ are i.i.d., we can bound the RHS by

$$\exp\Big[t\mathbb{E}\Big(e^{A|\xi_1|(t\vee 1)^{-1/2}} - 1\Big)\Big] \leq \exp\left(\sum_{k=1}^{m} r_k \frac{\cosh\big[kA(t\vee 1)^{-1/2}\big] - 1}{[(t\vee 1)^{-1/2}]^2}\right) \leq C.$$

Since $T \leq \tilde{T}$, this proves (A.10). The bound (A.11) follows from the preceding inequality and the fact that $\sup_{\mathbb{R}} |\zeta|^u e^{-\zeta} < \infty$. ☐

Lemma A.2. *For any $A \geq 0$, $v \in [0,1]$, $|j'|, |j''| \leq m$, $1 \leq j \leq k \leq m$, $|l| < m$,*

$$\begin{aligned} \varepsilon \sum_X \big|\nabla^- \mathbf{P}^\varepsilon_T(X - j'\varepsilon)\nabla^- \mathbf{P}^\varepsilon_T(X - j''\varepsilon)\big| e^{A|X|} &\leq \varepsilon^2 \big(T \vee \varepsilon^{-2}\big)^{-\frac{3}{2}} C, \\ \sum_x \big|\nabla^- \mathbf{p}_t(x - j')\nabla^- \mathbf{p}_t(x - j'')\big| e^{A\varepsilon|x|} &\leq \Big(t^{-\frac{3}{2}} \wedge 1\Big) C, \end{aligned} \tag{A.12}$$

$$\sum_x |(\mathbf{q}_{jkl})_t(x)| e^{A\varepsilon|x|} \leq \Big(1 \wedge t^{-\frac{3}{2}}\Big) C, \tag{A.13}$$

$$\sum_x |(\mathbf{q}_{jkl})_t(x)| |\varepsilon x|^v e^{A\varepsilon|x|} \leq \Big[\big(\varepsilon^2(t\vee 1)\big)^{\frac{v}{2}} (t\vee 1)^{-3/2} + \varepsilon^v (t\vee 1)^{-1}\Big] C. \tag{A.14}$$

Proof. We first prove (A.12) using the macroscopic variables. By Cauchy-Schwartz inequality the LHS of (A.12) can be bounded by a constant multiple of

$$\varepsilon \sum_X \big(\nabla^- \mathbf{P}^\varepsilon_T(X)\big)^2 e^{A|X|}. \tag{A.15}$$

Next, we apply discrete integration by parts to turn (A.15) into

$$-\varepsilon \sum_X \Big(\Delta \mathbf{P}^\varepsilon_T(X) \mathbf{P}^\varepsilon_T(X) e^{A|X|} + \nabla^- \mathbf{P}^\varepsilon_T(X) \mathbf{P}^\varepsilon_T(X) \nabla^- e^{A|X|}\Big),$$

and use (A.5), (A.7), and (A.10) to conclude (A.12). Note that $(T \vee \varepsilon^{-2})^{-1} \leq (T \vee \varepsilon^{-2})^{-3/2} C$, for $T \leq \tilde{T}$.

The bound (A.13) follows directly from (A.12). Next, using Cauchy-Schwartz inequality and discrete integration by parts as before, we bound the LHS of (A.14) by a constant multiple of

$$\sum_x \Big(|\Delta \mathbf{p}_t(x)| \mathbf{p}_t(x) |\varepsilon x|^v e^{A\varepsilon|x|} + \big|\nabla^- \mathbf{p}_t(x)\big| \mathbf{p}_t(x) (\varepsilon^v + \varepsilon|\varepsilon x|^v) e^{A\varepsilon|x|}\Big).$$

Using (A.5), (A.7), and (A.11), we conclude (A.14). ☐

Proof of Lemma 5.1. First, using discrete integration by parts we have

$$\sum_x \nabla^- \mathbf{p}_t(x-k) \nabla^- \mathbf{p}_t(x-l) = -\sum_x \mathbf{p}_t(x-k) \Delta \mathbf{p}_t(x-l) = -\sum_x \Delta \mathbf{p}_t(x-k+l) \mathbf{p}_t(x).$$

We then apply this identity and the telescoping identity (2.9) to the definition (5.3) of $\rho_{|k-l|}$ to obtain

$$\sum_{|l'|<k'} r_{k'}(k' - |l'|)\rho_{|k-l'|} = -r_{k'} \int_0^\infty \sum_x \mathbf{p}_t(x-k)\, \Delta_{k'} \mathbf{p}_t(x) dt, \tag{A.16a}$$

$$\sum_{|l'|<k'} r_{k'}(k' - |l'|)\rho_{|k-l'|} = -r_{k'} \int_0^\infty \sum_x \left(\Delta_{k'} \mathbf{p}_t(x-k)\right) \mathbf{p}_t(x) dt. \tag{A.16b}$$

For (5.5), we sum the identities (A.16) over k' and use (3.1) to get

$$\sum_{k'=1}^{m} \sum_{|l|'<k'} r_{k'}(k' - |l'|)\rho_{|k-l'|} = -\int_0^\infty \partial_t \Bigg(\sum_x \mathbf{p}_t(x)\mathbf{p}_t(x+k)\Bigg) dt,$$

which vanishes since, for $k \neq 0$, $\sum_x \mathbf{p}_t(x)\mathbf{p}_t(x+k) = 0$ at $t = 0$ and $t = \infty$.

For (5.6), first note that from discrete integration by parts and Cauchy-Schwartz inequality we have

$$\Big|\sum_x \mathbf{p}_t(x-k)\,\Delta_{k'}\mathbf{p}_t(x)\Big| < \sum_x \big(\nabla^-_{k'}\mathbf{p}_t(x)\big)^2, \tag{A.17}$$

where the inequality is strict with $k \neq 0$, since for any probability kernel $p(x)$ its discrete gradient $\nabla^-_{k'}p(x)$ cannot be periodic in x. Applying again discrete integration by parts to the RHS we obtain

$$\sum_x \big(\nabla^-_{k'}\mathbf{p}_t(x)\big)^2 = -\sum_x (\Delta_{k'}\mathbf{p}_t(x))\mathbf{p}_t(x). \tag{A.18}$$

Combining (A.16a), (A.17), (A.18), and (3.1), we obtain

$$\sum_{k'=1}^{m} \Big|\sum_{|l'|<k'} r_{k'}(k'-|l'|)\rho_{|k-l'|}\Big| < -\int_0^\infty \partial_t\Big(\sum_x \mathbf{p}_t(x)\mathbf{p}_t(x)\Big)dt = 1.$$

□

Proposition A.3. *Define a square matrix* $\hat\rho$ *of dimension* $2m-1$ *as*

$$\hat\rho_{ll'} = \int_0^\infty \sum_x |(\mathbf{q}_{ll'})_t(x)|dt, \quad |l|,|l'| < m.$$

For any $m \geq 2$*, there exists some* $(r_1,\ldots,r_m)$ *satisfying* (1.7) *and* (1.9) *such that the spectrum radius of* $\hat\rho$ *is strictly greater than one.*

Proof. In view of Gelfand's formula $\sigma(\hat\rho) = \lim_{n\to\infty}\|\hat\rho^n\|^{1/n}$ and the fact that all entries of $\hat\rho$ are non-negative, we need only to show that $\hat\rho_{ll} > 1$ for some $|l| < m$. Further, since by definition $\hat\rho_{ll'} \geq |\rho_{ll'}|$, it suffices to show that $|\rho_{ll}| > 1$ for some $|l| < m$.

From (5.4), we have

$$\rho_{m-1\,m-1} = r_m \int_0^\infty \sum_x \nabla^-\mathbf{p}_t(x-1)\nabla^-\mathbf{p}_t(x)dt.$$

Further, we use (A.1) and Parseval's identity to get

$$\rho_{m-1\,m-1} = \frac{r_m}{\pi}\int_{-\pi}^{\pi} \frac{\sin^2\left(\frac{\zeta}{2}\right)\cos\zeta}{\sum_{k=1}^m r_k(1-\cos(k\zeta))}d\zeta.$$

Now, set $(r_1,\ldots,r_m) = (\delta,0,\ldots,0,1-\delta)$ with $\delta \in (0,1)$, which satisfies (1.7) and (1.9). Since $1-\cos(m\zeta)$ vanishes at $\zeta_j := \frac{2j\pi}{m}$, elementary analysis using the following readily verified identity

$$\lim_{a\downarrow 0} a^{\frac12}\int_{-A}^{A}\frac{1}{a+\frac12(m\zeta)^2}d\zeta = \frac{\pi}{m}\left(\frac{2}{a}\right)^{\frac12}, \text{ for any } A > 0$$

shows that

$$\lim_{\delta\downarrow 0}\delta^{\frac12}\rho_{m-1\,m-1}(\delta) = \frac{2^{\frac12}\pi}{m}\left(\sum_{0<|j|<\frac{m}{2}} w(\zeta_j) + \mathbb{1}_{\{m/2\in\mathbb{N}\}}w(\zeta_{m/2})\right), \tag{A.19}$$

where

$$w(x) := \frac{\sin^2(x/2)\cos x}{(1-\cos x)^{\frac12}} = (1-\cos x)^{\frac12}\cos x.$$

Since $1-\cos x$ is increasing over $[0,\pi]$, by considering separately the cases $m/2 \in \mathbb{N}$ and $m/2 \notin \mathbb{N}$, we know that the RHS of (A.19) is negative when $m > 1$, which implies that $\rho_{m-1\;m-1} \to -\infty$ as $\delta \to 0$. □

References

[1] G. Amir, I. Corwin, and J. Quastel. Probability distribution of the free energy of the continuum directed random polymer in 1 + 1 dimensions. *Comm. Pure Appl. Math.*, 64(4):466–537, 2011.

[2] L. Bertini and N. Cancrini. The stochastic heat equation: Feynman-Kac formula and intermittence. *J. Stat. Phys.*, 78(5-6):1377–1401, Mar. 1995.

[3] L. Bertini and G. Giacomin. Stochastic burgers and KPZ equations from particle systems. *Comm. Math. Phys.*, 183(3):571–607, Feb. 1997.

[4] A. Borodin and I. Corwin. Macdonald processes. *arXiv:1111.4408*, Nov. 2011.

[5] A. Borodin, I. Corwin, and P. Ferrari. Free energy fluctuations for directed polymers in random media in 1+1 dimension. *arXiv:1204.1024*, Apr. 2012.

[6] I. Corwin. The Karder-Parisi-Zhang equation and universality class. *Random Matrices: Theory Appl.*, 01(01):1130001, Jan. 2012.

[7] I. Corwin and J. Quastel. Crossover distributions at the edge of the rarefaction fan. *arXiv:1006.1338*, June 2010.

[8] D. Forster, D. R. Nelson, and M. J. Stephen. Large-distance and long-time properties of a randomly stirred fluid. *Physical Review A*, 16(2):732–749, Aug. 1977.

[9] A. Friedman. *Partial Differential Equations of Parabolic Type*. Dover Publications, Apr. 2008.

[10] J. Gärtner. Convergence towards Burger's equation and propagation of chaos for weakly asymmetric exclusion processes. *Stochastic Process. Appl.*, 27(0):233–260, 1987.

[11] M. Hairer. Solving the KPZ equation. *arXiv:1109.6811*, Sept. 2011.

[12] D. A. Huse, C. L. Henley, and D. S. Fisher. Huse, henley, and fisher respond. *Phys. Rev. Lett.*, 55(26):2924–2924, Dec. 1985.

[13] K. Johansson. Shape fluctuations and random matrices. *Comm. Math. Phys.*, 209(2):437–476, Feb. 2000.

[14] K. Johansson. Discrete polynuclear growth and determinantal processes. *Comm. Math. Phys.*, 242(1-2):277–329, Nov. 2003.

[15] M. Kardar, G. Parisi, and Y.-C. Zhang. Dynamic scaling of growing interfaces. *Phys. Rev. Lett.*, 56(9):889–892, Mar. 1986.

[16] C. Mueller. On the support of solutions to the heat equation with noise. *Stochastics*, 37(4):225–245, 1991.

[17] M. Prähofer and H. Spohn. Scale invariance of the PNG droplet and the Airy process. *J. Stat. Phys.*, 108(5-6):1071–1106, Sept. 2002.

[18] T. Sasamoto and H. Spohn. Exact height distributions for the KPZ equation with narrow wedge initial condition. *Nucl. Phys. B*, 834(3):523–542, Aug. 2010.

[19] C. A. Tracy and H. Widom. A Fredholm determinant representation in ASEP. *J. Stat. Phys.*, 132(2):291–300, July 2008.

[20] C. A. Tracy and H. Widom. Integral formulas for the asymmetric simple exclusion process. *Comm. Math. Phys.*, 279(3):815–844, May 2008.

[21] C. A. Tracy and H. Widom. Asymptotics in ASEP with step initial condition. *Comm. Math. Phys.*, 290(1):129–154, Aug. 2009.

[22] C. A. Tracy and H. Widom. On ASEP with step bernoulli initial condition. *J. Stat. Phys.*, 137(5-6):825–838, Dec. 2009.

[23] C. A. Tracy and H. Widom. Formulas for ASEP with two-sided Bernoulli initial condition. *J. Stat. Phys.*, 140(4):619–634, Aug. 2010.

[24] J. B. Walsh. An introduction to stochastic partial differential equations. In P. L. Hennequin, editor, *École d'Été de Probabilités de Saint Flour XIV - 1984*, number 1180 in Lecture Notes in Mathematics, pages 265–439. Springer Berlin Heidelberg, Jan. 1986.

*† Department of Mathematics, Stanford University, California 94305